



Management of *Rhizoctonia solani*, *Fusarium solani* and *Meloidogyne incognita* by Silicon dioxide nanoparticles and *Rhizobium ciceri* alone and in combination on chickpea

Zaki A Siddiqui* and Sumaiya Aziz

Department of Botany, Aligarh Muslim University, India

*Corresponding author: Zaki A Siddiqui, Department of Botany, Aligarh Muslim University, India

Received: 📅 September 08,2020

Published: 📅 September 23,2020

Abstract

Effects of silicon dioxide nanoparticles (SiO₂ NPs) and *Rhizobium ciceri* was observed alone and in combination on *Rhizoctonia solani* / *Fusarium solani* / *Meloidogyne incognita* on the growth, photosynthetic pigments (chlorophyll and carotenoid) and proline contents of *Cicer arietinum*.

Inoculation of *M. incognita* resulted in a greater reduction in plant growth, photosynthetic pigments and higher increase in proline contents than other test pathogens. Inoculation of *R. ciceri* or foliar spray of SiO₂ NPs (0.10 mg ml⁻¹) with pathogens under study resulted in increased plant growth, photosynthetic pigments and proline contents than without *R. ciceri* / NPs. Combined application of NPs with *R. ciceri* resulted in a greater increase in plant growth, photosynthetic pigments and proline contents in plants with pathogens than with NPs or *R. ciceri*. Plants without *R. ciceri* had a very poor root nodulation but nodulation was high in plants with *R. ciceri*. Pathogens under study and NPs had adverse effect on nodulation caused by *R. ciceri*. Wet rot and black root rot indices were 4 when *R. solani* and *F. solani* were inoculated respectively. *R. ciceri* / NPs reduced wet root rot, black root rot indices, galling and population of *M. incognita*. Disease indices, galling and population of *M. incognita* were reduced greatly by use of *R. ciceri* plus SiO₂ NPs.

Keywords: Black Root Rot; *Cicer Arietinum*; Disease Index; Root Knot Nematode; Root Nodule Bacterium; Wet Root Rot

Introduction

Chickpea (*Cicer arietinum* L.) is widely grown as major food legume because of its rich nutrient values. It has high protein and starch suitable for textile sizing Duke [6]. *Rhizobium* Frank is root nodule bacterium associated with roots of legumes for symbiotic nitrogen fixation. *Rhizobium* inoculation has positive effect on growth attributes, yield components and quality of chickpea. Application of rhizobial isolates significantly suppress root rotting fungi and root knot nematode Parveen et al. [20] and reduced damaged caused Siddiqui and Mahmood [23].

Important diseases of chickpea include; *Rhizoctonia* wet rot (*Rhizoctonia solani* Kuhn) and *Fusarium* black root rot (*Fusarium solani* Mart. Amend Sacc.) and root-knot disease by *Meloidogyne*

incognita (Kofoid and White) Chitwood. Parasitism by *M. incognita* in chickpea involves giant cells formation and root galling Vovlas et al. [29]. *Rhizoctonia* wet root causes severe losses from seedling to maturity stages Jayalakshmi et al. [12]. Its characteristic symptoms include root rotting and gradual yellowing and wilting of foliage, the rotted, discolored tissues are soft and wet. *Fusarium* black root rot also results in severe losses to of chickpea crop Jayalakshmi et al. [12]. The root system is generally rotted with most of the finer roots destroyed, while the taproot remains intact, but dark and necrotic.

Silicon (Si) application is useful for increasing plant growth. Use of Si overcome biotic stresses and also improves plant's

physiological and mechanical properties Epstein [7]; Ma and Yamaji [10]. Si also enhances disease resistance against pest and pathogens Marschner [18].

Frequent occurrence of *F. solani*, *R. solani* and *M. incognita* was observed in the chickpea fields of Aligarh, U.P. India. Infected plants were found with symptoms of diseases having poor plant growth. Silicon dioxide nanoparticles (SiO₂ NPs) and *Rhizobium ciceri* were used alone and in combination management of *F. solani*, *R. solani* and *M. incognita* on chickpea. Effects of SiO₂ NPs and *R. ciceri* were also observed on chickpea growth, photosynthetic pigments and also on proline contents.

Materials And Methods

Chickpea fields were surveyed for the collection disease materials of Aligarh U.P. Samples (root and soil) were collected and placed at 4°C until processing in refrigerator. The presence of plant parasitic nematodes and fungi in the collected samples were examined.

Isolation of fungi from infected chickpea

Roots showing disease symptoms were washed with sterilized water three times to remove soil particles. Isolation of fungi was made from infected plant parts Siddiqui et al. [24] on potato dextrose agar (PDA) medium. Identity of *Rhizoctonia solani* and *Fusarium solani* was confirmed. Later, pure cultures were separately prepared of both *R. solani* and *F. solani*

Root-knot nematode *M. incognita*

Root-knot nematodes females were dissected out from roots of chickpea having symptoms of root-knot. Later, perineal patterns were prepared for the identification of *Meloidogyne* sp. Taylor and sasser [27].

Preparation of SiO₂ NPs suspension

SiO₂ NPs (Sigma-Aldrich, product No. 637246-50G) was used to prepare 0.10 mg ml⁻¹ suspension by dissolving 100 mg nanopowder in 1-liter sterilized water. Foliar spray of 10 ml suspension was made on seedlings (15-day old) in a greenhouse with small spray pumps. Distilled water (10 ml) was sprayed on control plants and each treatment had five replications.

Effect of SiO₂ NPs on *M. incognita*

The effect of SiO₂ NPs was studied on the *M. incognita*. Egg masses (20) of average size from eggplant roots were picked with sterilized forceps and effect of 0.10 mg ml⁻¹ NPs on the hatching and mortality was observed as described by Siddiqui et al. [24].

Effect of SiO₂ NPs on the fungi

Activity of SiO₂ NPs against both fungi was observed separately. Ten ml NPs (0.10 mg ml⁻¹ conc.) was added in 100 ml autoclaved PDA to observe effect as described by Siddiqui et al. [24]

Preparation and sterilization of soil mixture

Collected loam soil was sieved through 10 meshes. Clay pots were filled with 1 kg of soil mixed with loam soil and river sand (3:1). Before sterilization, soil surface of pots was wet with water, sterilized at 137.9 kPa for 20 minutes and allowed for cooling 24 hours.

Raising and maintenance of test plants

Chickpea (*Cicer arietinum* L.) seeds Pusa-3022 were sterilized and sown in 15 cm pots Siddiqui et al. [24]. Seedlings were inoculated and un-inoculated control were placed in a glass house at 20 ± 2°C (Table 2). Pots were arranged on a bench and each treatment had five replications. Watering of plants was done as required. Ninety days after inoculation plants were harvested

Inocula of fungi

Richard's liquid medium was used for culturing of both fungi separately Riker and Riker [22] for obtaining inocula. Eighty ml liquid medium after filtration through muslin cloth was placed 250 ml flasks, autoclaved at 103.4 kPa for 15 minutes. Each fungus was separately inoculated, incubated, mycelia mats were collected, blotting sheets were used to remove excess water and inoculum of each fungus was prepared in Waring blender as described Siddiqui et al. [24]. Ten ml of this suspension was used as inoculum (1 g fungus).

Nematode inoculum

Infected chickpea roots were used to collect *Meloidogyne incognita*, eggplants for its multiplication. Later, inoculum was prepared, juveniles were counted and 2000 juveniles were inoculated as described by Siddiqui et al. [24]

Rhizobium inoculum

Rhizobium ciceri (100 g) charcoal culture was dissolved in 1 liter sterilized water. Ten ml (1 g inoculum) around the seeds per pot was inoculated where required.

Inoculation technique

Rhizobium ciceri was used with seeds at sowing in about half seeds. Inoculation of *R. solani*, *F. solani* and *M. incognita*, chickpea seedlings (two weeks old) were used and inoculations were done Siddiqui et al. [24]. There were 2 treatments i.e. (1) Control; (2) *Rhizobium ciceri*. Each of these was tested with (I) Control; (II) *R. solani*; (III) *F. solani*; (IV) *M. incognita* (4×2=8). These 8 treatments were tested in two combinations (i) without SiO₂ NPS spray (ii) With SiO₂ NPs spray (8×2= 16). Each treatment had 5 replications (16×5=80 pots).

Evaluation of the experiment

Ninety days after inoculation the plants were harvested. Plant growth attributes, photosynthetic pigments, proline and disease

index, galling and nematode population were recorded Siddiqui et al., [24].

Disease index

Disease index were determined on disease symptoms. Disease rating was done on 0 (no disease) to 5 (severe rot / blight) Nesha and Siddiqui [19]

Estimation of photosynthetic pigments and proline content

Photosynthetic pigments (chlorophyll and carotenoid) in the fresh leaf samples were estimated by Mackinney [17] and proline content by Bates et al. [3].

Statistical analysis

The data were analysed through three ways (*Rhizobium ciceri* × SiO₂ NPs × Pathogens) analysis of variance (ANOVA) in R (version 2.14.0) statistical software (package- library agricolae). Least

significant difference (L.S.D) were calculated at p=0.05. Duncan's new multiple range test (DNMRT) were employed to denote the significant differences between treatments. Graphs of nematode population and galls per root system were prepared using Sigma Plot™ and error bars showing standard error.

Results

Effects on SiO₂ NPS on pathogenic fungi and on *M. incognita*

Effects of SiO₂ NPs were studied on *R. solani* and *F. solani* (Table 1). It had a higher adverse effect on the growth of *F. solani* followed by *R. solani*. SiO₂ NPs caused 36.25 and 34.60% inhibitions in the growth of *F. solani* and *R. solani* respectively (Table 1). SiO₂ NPs also had adverse effect on the hatching of *M. incognita*. It caused 88.37% inhibition in hatching of *M. incognita* while 43.75% mortality of *M. incognita* J2 was observed after 48 hours of incubation over control (Table 1).

Table 1: Effect of SiO₂ NPs on the growth of *F. solani* and *R. solani* in potato dextrose agar medium and on the hatching and mortality of *M. incognita*.

Nanoparticles	Concentration	Fusarium solani	
		Colony diameter (cm)	% inhibition over control
Control	DDW	8.00 a	-
SiO ₂ NPs	0.10 mg ml ⁻¹	5.10 b	36.25
L.S.D. P≤0.05		0.44	-
Nanoparticles	Concentration	Hatching of <i>M. incognita</i> J2	% inhibition over control
Control	DDW	215 a	-
SiO ₂ NPs	0.10 mg ml ⁻¹	19 b	88.37

Pot experiment

Three ways ANOVA revealed that individual effect of *Rhizobium ciceri*, SiO₂ NPs and pathogen, and interaction of *R. ciceri* × SiO₂ NPs, Pathogen × *R. ciceri*, SiO₂ NPs × pathogen and *R. ciceri* × SiO₂ NPs × pathogen was significant in proline content and root nodulation at p=0.05 (ANOVA not shown). However, effect of interaction of *R. ciceri* × SiO₂ NPs × pathogen on shoot length, root fresh weight, root dry weight and photosynthetic pigments was non-significant at p=0.05. Similarly, effect of interaction of Pathogen × *R. ciceri* on root length, shoot fresh weight, shoot dry weight, root dry weight, chlorophyll was also non-significant at p=0.05. Effect of interaction on SiO₂ NPs × pathogen on shoot fresh weight, root dry weight and photosynthetic pigments were also non-significant at p=0.05. However, Individual effect of *R. ciceri*, SiO₂ NPs and pathogen and their interactions were significant on other parameters studied (ANOVA not shown).

Effect on plant growth and nodulation

Inoculation of plants with pathogens under study i. e. *M. incognita*, *F. solani* and *R. solani* resulted in a significant reduction in plant growth parameters over uninoculated control (Table 2).

Inoculation of *M. incognita* resulted in a higher reduction (26.07%) in shoot dry weight followed by *R. solani* (23.92%) and *F. solani* (20.10%). Similarly, inoculation of *M. incognita* resulted in a higher reduction (25.42%) in root dry weight followed by *R. solani* (23.73%) and *F. solani* (20.33%). Plants without *R. ciceri* had poor nodulation while plants with *R. ciceri* had higher nodulation (Table 2).

Foliar spray of SiO₂ NPs resulted in a significant increase in plant growth parameters over plants with and without pathogen (Table 2). Spray of SiO₂ NPs to plants without pathogen caused 25.36% increase in shoot dry weight over uninoculated control. Foliar spray of SiO₂ NPs to plants with *M. incognita* caused 33.01% increase in shoot dry weight while plants with *F. solani* and *R. solani* resulted in 39.22% and 39.31% increase in shoot dry weight over plants with respective pathogen alone (Table 3). Foliar spray of SiO₂ NPs caused 38.98% increase in root dry weight over uninoculated control. Spray of SiO₂ NPs to plants with *M. incognita*, *F. solani* and *R. solani* resulted in 45.45%, 53.19% and 53.33% increase in root dry weight respectively. Nodulation in plants without *R. ciceri* remain unaffected by inoculation of pathogens and SiO₂ NPs spray because of very poor nodulation (Table 2).

Table 2: Effects of SiO₂ NPs and *Rhizobium ciceri* alone and both together on *F. solani*, *R. solani* and *M. incognita* and on the growth attributes and nodulation of chickpea. Values within a column followed by the same letter are not significantly different at p=0.05 by DNMRT.

Treatment		Shoot length (cm)	Root length (cm)	Shoot fresh weight (g)	Root fresh weight (g)	Shoot dry weight (g)	Root dry weight (g)	No of nodules / root system	
Nil	Nil	C	48.6cd	16.0gh	14.37h	6.90l	4.18hi	0.59g	4g
		M	35.8i	12.2l	11.01j	4.93n	3.09k	0.44h	3gh
		F	40.0h	13.9j	12.31i	5.29m	3.34j	0.47h	3gh
		R	38.2h	12.7k	11.39j	5.20m	3.18jk	0.45h	2h
	SiO ₂ NPs Spray	C	51.2ab	18.4d	16.65f	9.08g	5.24d	0.82cd	3gh
		M	44.4g	15.2i	14.19h	7.19k	4.11i	0.64fg	4g
		F	46.0efg	16.3g	15.09g	7.92i	4.65f	0.72def	3gh
		R	45.2fg	15.8h	14.49h	7.50j	4.43fg	0.69efg	3gh
Rhizobium ciceri	Nil	C	50.4bc	18.8cd	20.20c	10.77e	5.26d	0.85c	46a
		M	44.2g	16.0gh	17.52e	8.63h	4.35gh	0.70ef	36c
		F	45.8efg	17.8e	18.40d	9.57f	5.02e	0.79cde	32d
		R	44.6g	16.8f	18.13d	9.21g	4.62f	0.75cde	29e
	SiO ₂ NPs Spray	C	53.0a	21.8a	22.93a	13.18a	8.77a	1.30a	42b
		M	47.2def	18.7cd	19.97c	11.19d	7.29c	1.05b	31d
		F	48.2cde	19.3b	20.92b	12.35b	7.92b	1.14b	28e
		R	47.8de	18.9bc	20.05c	12.08c	7.44c	1.09b	24f
LSD p=0.05		2.21	0.45	0.41	0.24	0.21	0.09	1.67	

Table 3: Effects of SiO₂ NPs and *Rhizobium ciceri* alone and both together on *F. solani*, *R. solani* and *M. incognita* and on the photosynthetic pigments (chlorophyll and carotenoid) proline content and disease index of chickpea.

Treatments		Total chlorophyll (mg/FW)	Carotenoids content (mg/FW)	Proline content (µmol/g FW)	Root rot disease index	
Nil	Nil	C	0.198g	0.0675h	0.0024l	-
		M	0.136i	0.0501j	0.0052jk	-
		F	0.186gh	0.0640hi	0.0028l	4
		R	0.174h	0.0606i	0.0047k	4
	SiO ₂ NPs Spray	C	0.236de	0.0880ef	0.0104g	-
		M	0.201fg	0.0687h	0.0171e	-
		F	0.226e	0.0860f	0.0107g	3
		R	0.219ef	0.0783g	0.0139f	3
Rhizobium ciceri	Nil	C	0.259c	0.1002c	0.0061ij	-
		M	0.237de	0.0893def	0.0094gh	-
		F	0.257c	0.0940d	0.0068i	3
		R	0.248cd	0.0917de	0.0083h	3
	SiO ₂ NPs Spray	C	0.288a	0.1127a	0.0184d	-
		M	0.264bc	0.1022bc	0.0331a	-
		F	0.286a	0.1059b	0.0234c	2
		R	0.283ab	0.1041bc	0.0268b	2
LSD p=0.05		0.020	0.005	0.001	-	

Values within a column followed by the same letter are not significantly different at p=0.05 by DNMRT.

Inoculation of *R. ciceri* resulted in a significant increase in plant growth parameters over pathogen inoculated and un-inoculated control (except shoot length in uninoculated plants) (Table 2). Inoculation of *R. ciceri* resulted in 25.84% increase in shoot dry weight over uninoculated control. Use of *R. ciceri* to plants with *M. incognita*, *F. solani* and *R. solani* resulted in 40.78, 50.30 and 45.28% increase in shoot dry weight respectively over their respective inoculated control (Table 2). Inoculation of *R. ciceri* caused 44.06% increase in root dry weight over uninoculated control. Similarly, chickpea plants grown with *R. ciceri* and with *M. incognita*, *F. solani* and *R. solani* resulted in 59.09, 68.09 and 66.67% increases in root dry weight over respective pathogen inoculated plants. High root nodulation was observed in plants with *R. ciceri* while pathogens under study had adverse effect on root nodulation caused by *R. ciceri* (Table 2).

Foliar spray of SiO₂ NPs together with *R. ciceri* caused a highest increase in plant growth parameters over both un-inoculated and pathogen inoculated control (Table 2). Spray of SiO₂ NPs with *R. ciceri* resulted in 109.80% increase in shoot dry weight over control. Similarly, spray of SiO₂ NPs plus inoculation of *R. ciceri* to plants with *M. incognita*, *F. solani* and *R. solani* caused 139.92, 137.13 and 133.96% increase in shoot dry weight over their respective inoculated control (Table 2). Use of *R. ciceri* with SiO₂ NPs resulted in 120.33% increase in root dry weight over control. However, spray of SiO₂ NPs to plants with *R. ciceri* and test pathogens i.e. *M. incognita*, *F. solani* and *R. solani* resulted in 138.64, 142.55 and 142.22% increase in root dry weight over respective control. Pathogen under study had adverse effect on root nodulation caused by with *R. ciceri* and spray of SiO₂ NPs also adversely affect root nodulation (Table 2).

Effect on photosynthetic pigments

Inoculation of plants with *M. incognita*, and *R. solani* caused a significant reduction in photosynthetic pigments i. e. chlorophyll and carotenoid contents over uninoculated plants (Table 3). Foliar spray of SiO₂ to plants with pathogens under study caused a significant increase in photosynthetic pigments over plants with pathogen alone. Foliar application of SiO₂ caused 19.19 and 30.37%

increases in chlorophyll and carotenoid respectively over control. Similarly, foliar spray of SiO₂ to plants with *M. incognita*, *R. solani* and *F. solani* resulted in 21.50 to 47.79% increase in chlorophyll and 29.21 to 37.13% increase in carotenoid over respective inoculated control. Inoculation of *R. ciceri* caused 30.80 and 48.44% increase in chlorophyll and carotenoid respectively over control. Similarly, *R. ciceri* inoculation to plants with *M. incognita*, *R. solani* and *F. solani* resulted in 42.53 to 74.26% increase in chlorophyll and 46.88 to 78.24% increase in carotenoid contents. Combined use of *R. ciceri* together with SiO₂ NPs caused 45.45 and 66.96% increase in chlorophyll and carotenoid respectively over control. Similarly, use of *R. ciceri* with SiO₂ NPs to plants with *M. incognita*, *R. solani* and *F. solani* caused 53.76 to 94.11% increase in chlorophyll and 65.47 to 103.99% increase in carotenoid contents (Table 3).

Effect on proline contents

A significant increase in proline contents was observed in plants with *M. incognita*, *F. solani* and *R. solani* over control (Table 3). Plants with *M. incognita*, caused a higher increase in proline contents than by *F. solani*. Foliar spray of SiO₂ or inoculation of *R. ciceri* also increased proline contents. Use of SiO₂ NPs together with *R. ciceri* resulted in highest increase in proline contents over other treatments (Table 3).

Disease indices

Wet root rot and black root rot disease indices were found 4 in plants with *R. solani* and *F. solani* respectively (Table 3). Disease indices were 3 in plants with *R. solani* / *F. solani* and sprayed with SiO₂ NPs or inoculated with *R. ciceri*. Combined use of SiO₂ NPs with *R. ciceri* reduced disease indices to 2 (Table 3).

Root galling and nematode population

Inoculation of *M. incognita* inoculated alone resulted in high root galling and nematode population (Figure 1, 2). Foliar spray of SiO₂ NPs or inoculation of *R. ciceri* reduced galling and nematode population. SiO₂ NPs spray caused a higher reduction in galling and nematode population as compared to *R. ciceri* inoculation. Spray of SiO₂ NPs together with *R. ciceri* caused highest reduction in nematode population and galling (Figure 1, 2).




Figure 1: Effect of SiO₂ NPs (NPs) and *Rhizobium ciceri* (R) on the population of *M. incognita* (M) on chickpea.




Figure 2: Effect of SiO₂ NPs (NPs) and *Rhizobium ciceri* (R) on the galling of *M. incognita* (M) on chickpea.

Discussion

SiO₂ NPs showed antifungal effect in PDA medium because growth of *R. solani*, and *F. solani* was inhibited. Similarly, Akpınar et al. [1] found that SiO₂ NPs resulted in maximum inhibition of *F. oxysporum* f. sp. *radicis lycopersici*. SiO₂ NPs also showed adverse effect on *M. incognita* J2 hatching. Observation of hatched J2 showed straight body shape and intake of NPs in J2 of *M. incognita* within 48 h of incubation. Silica nanoparticles caused degeneration of reproductive organs of nematodes Pluskota et al. [21]. The silicon carbide nanoparticles had lethality as dead nematodes exhibited black internal organs Al Banna et al. [2]. It is possible that SiO₂ NPs taken by the nematodes were translocated to primary and

secondary organs Wu et al. [32]; Al Banna et al. [2] resulting in deformation of J2 of *M. incognita*.

In pot experiment, foliar spray of SiO₂ NPs increased growth, chlorophyll and carotenoids contents over uninoculated control. Application of Si in plants results in its accumulation and exerts various beneficial effects on plants Liang et al. [14]. Absorbed Si alleviates various abiotic stresses Wang et al. [31]. Si contributes to overcome stresses also may improves plant's mechanical and physiological properties Epstein [7]; Ma and Yamaji [16]. Increased growth of plants without pathogen (uninoculated control) by SiO₂ application can be due to above reasons.

Foliar spray of SiO₂ NPs increased growth attributes of plants with pathogen. Applications of Si provide resistance against diseases Fauteux et al. [8]; Marschner [18]. Various mechanisms are involved plant disease resistance i.e. defense barriers after pathogen infection, activation enzymes (defense-related), production of antimicrobial compounds and regulation of genes Wang et al. [31]. Si mediated resistance against pathogens was associated with the changes in activity of soil enzyme and soil microorganisms Wang et al. [31]. Silicon amendment also reduced disease incidence of tomato Diogo and Wydra [5].

Foliar spray of SiO₂ can prevent pathogen penetration by formation of a cuticle-Si double layer and decrease disease incidence Ma and Yamaji [16]. Most Si improves mechanical properties and regeneration by cross-linking with hemicellulose in cell walls Guerriero et al. [11].

Proline contents were found to be increased by the spray of SiO₂ NPs in this study. Proline is known to confer tolerance to plants Lehmann et al. [13]. It is a multi-functional amino acid and accumulation of proline acts as a major stress response against pathogens Grote et al. [10]. A correlation between increase in proline contents and increase defense response against pathogens Cecchini et al. [4] has been observed. Increased proline content was also observed in Arabidopsis thaliana Verslues and Sharma [28] in plant defense against pathogen. Therefore, in this study increase in proline contents in plants with pathogen was also observed.

Disease indices were found 4 in plants with fungal pathogens viz. *R. solani* and *F. solani*. Foliar spray of SiO₂ reduced disease indices to 2. Reduction in disease indices by application of SiO₂ also confirms antifungal nature of SiO₂. SiO₂ NPs demonstrated its nematicidal activity by reduction in galling and nematode population. *R. ciceri* also showed antifungal and nematicidal activity by reducing disease indices, galling and *M. incognita* population therefore, increase in plant growth. Rhizobial strains have biocontrol properties and can lead to potential control Gopalakrishnan et al. [9]. *Rhizobium* enhanced resistance level against pathogen by inducing changes in seed proteome and metabolome Sistani et al. [26]. Foliar application of SiO₂ NPs had mild adverse effect on rhizobial root nodulation while combined application of NPs and *R. ciceri* reduce disease severity. This may be possible because of least concentration NPs used as foliar spray. The influence of broad-spectrum of even pesticides and their tolerance to the rhizobia are reported Singh and Wright [25]. It is possible that the *R. ciceri* used in the present study may have tolerance to SiO₂ NPs thereby resultant increase in growth attributes, chlorophyll, carotenoid and proline contents.

Conclusion

SiO₂ NPs had displayed antifungal and nematicidal activities towards tested fungi and root knot nematode. Similarly, *R. ciceri*

also provided biocontrol of *M. incognita* and root rot fungi. Use of SiO₂ NPs with *R. ciceri* was better for the management of pathogens under study than their use alone. Therefore, researches on characterizing of resistant rhizobia are needed so they can be used for the management of chickpea diseases along with SiO₂ NPs.

Acknowledgements

Necessary facilities provided by Aligarh Muslim University Aligarh, India are thankfully acknowledged.

References

- Akpinar I, Sar T, Unal M (2017) Antifungal effects of Silicon dioxide nanoparticles (SiO₂ NPs) against various plant pathogenic fungi. Conference: International Workshop Plant Health: Challenges and Solutions 23-28.
- Al-Banna L, Salem N, Ghrair A. M, Habash SS (2018) Impact of silicon carbide nanoparticles on hatching and survival of soil nematodes *Caenorhabditis elegans* and *Meloidogyne incognita*. Appl Ecol Env Res 16(3): 2651-2662.
- Bates LS, Waldren RT, Teare ID (1973) Rapid determination of free proline for water stress studies. Plant Soil 39: 205-207.
- Cecchini NM, Monteoliva MI, Alvarez ME (2011) Proline dehydrogenase contributes to pathogen defense in Arabidopsis. Plant Physiol 155(4): 1947-1959.
- Diogo RVC, Wydra K (2007) Silicon-induced basal resistance in tomato against *Ralstonia solanacearum* is related to modification of pectic cell wall polysaccharide structure. Physiol Mol Plant Pathol 70 (4-6): 120-129.
- Duke JA (1981) Handbook of legumes of world economic importance. Plenum Press New York 52-57.
- Epstein E (1999) Silicon. Annu Rev Plant Physiol Plant Mol Biol 50: 641-664.
- Fauteux F, Remus-Borel W, Menzies JG, Belanger RR (2005) Silicon and plant disease resistance against pathogenic fungi. FEMS Microbiol Lett 249(1): 1-6.
- Gopalakrishnan S, Sathya A, Vijayabharathi R, Varshney RK, Gowda CLL, et al. (2018) Plant growth promoting rhizobia: challenges and opportunities. 3 Biotech 5(4): 355-377.
- Grote D, Schmit R, Claussen W (2006) Water uptake and proline index as indicators of predisposition in tomato plants to *Phytophthora nicotianae* infection as influenced by abiotic stresses. Physiol Mol Plant Pathol 69(4-6): 121-130.
- Guerriero G, Hausman JF, Legay S (2016) Silicon and the plant extracellular matrix. Front Plant Sci 7: pp. 463.
- Jayalakshmi SK, Usharani S, Benagi VI, Mannur DM (2008) Sources of resistance to dry root rot of chickpea caused by *Rhizoctonia bataticola*. Agric Sci Digest 28 (2): 147 -148.
- Lehmann S, Funck D, Szabados L, Rentsch D (2010) Proline metabolism and transport in plant development. Amino Acids 39: 949-962.
- Liang YC, Wong JWC, Wei L (2005) Silicon-mediated enhancement of cadmium tolerance in maize (*Zea mays* L.) grown in cadmium contaminated soil Chemosphere 58(4): 475-483.
- Ma JF, Yamaji N (2006) Silicon uptake and accumulation in higher plants. Trends Plant Sci., 11(8): 392-397.
- Ma JF, Yamaji N (2008) Functions and transport of silicon in plants. Cell. Mol. Life Sci., 65(19): 3049-3057.

17. Mackinney G (1941) Absorption of light by chlorophyll solutions. J Biol Chem 140: 315-322.
18. Marschner P (2012) Marschner's Mineral Nutrition of Higher Plants. London: Academic Press pp 672.
19. Nesha R, Siddiqui ZA (2013) Interactions of *Pectobacterium carotovorum* pv. *carotovorum*, *Xanthomonas campestris* pv. *carotae*, and *Meloidogyne javanica* on the Disease Complex of Carrot. Int J Veg Sci 19(4): 403-411.
20. Parveen G, Noreen R, Shafique HA, Sultana V, Haque SE, et al. (2019) Role of rhizobia in suppressing the root diseases of soybean under soil amendment. Planta daninha 37.
21. Pluskota A, Horzowski E, Bossinger O, von Mikecz A (2009) In *Caenorhabditis elegans* nanoparticle-bio-interactions become transparent: silica-nanoparticles induce reproductive senescence. PLoS One 4(8): e6622.
22. Riker AJ, Riker RS (1936) Introduction to Research on Plant Diseases: A Guide to the Principles and Practice for Studying Various Plant-disease Problems. John's Swift Co. Inc., St. Louis, Chicago, New York, India 3: pp. 119.
23. Siddiqui ZA, Mahmood I (2001) Effects of rhizobacteria and root symbionts on the reproduction of *Meloidogyne javanica* and growth of chickpea. Bioresour Technol 79(1): 41-45.
24. Siddiqui ZA, Parveen A, Ahmad L, Hashem A (2019) Effects of graphene oxide and zinc oxide nanoparticles on growth, chlorophyll, carotenoids, proline contents and diseases of carrot. Sci Hort 249: 374-382.
25. Singh, G, Wright D (2002) In vitro studies on the effects of herbicides on the growth of rhizobia. Lett Appl Microbiol 35(1):12-16.
26. Sistani NR, Kaul HP, Desalegn G, Wienkoop S (2017) *Rhizobium* Impacts on Seed Productivity, Quality, and Protection of *Pisum sativum* upon Disease Stress Caused by *Didymella pinodes*: Phenotypic, Proteomic, and Metabolomic Traits. Front Plant Sci 8: 1961.
27. Taylor AL, Sasser JN (1978) Biology, Identification and Control of Root Knot Nematodes (*Meloidogyne* species). A cooperative publication of North Carolina State University, Dept of Plant Pathology and USAID, Raleigh NC, USA 7: pp. 111.
28. Verslues PE, Sharma S (2010) Proline metabolism and its implications for plant-environment interaction. Arabidopsis Book 8: e0140.
29. Vovlas N, Rapoport HF, Jiménez Díaz RM, Castillo P (2005) Differences in feeding sites induced by root-knot nematodes, *Meloidogyne* spp., in chickpea. Phytopathology 95(4): 368-375.
30. Wang L, Cai K, Chen Y, Wang G (2013) Silicon-mediated tomato resistance against *Ralstonia solanacearum* is associated with modification of soil microbial community structure and activity. Biol Trace Elem Res 152(2): 275-283.
31. Wang M, Gao L, Dong S, Sun Y, Guo SQ (2017) Role of Silicon on Plant-Pathogen Interactions. Front Plant Sci 8: pp. 701.
32. Wu S, Lu J, Rui Q, Yu S, Cai T, et al. (2011) Aluminum nanoparticle exposure in L1 larvae results in more severe lethality toxicity than in L4 larvae or young adults by strengthening the formation of stress response and intestinal lipofuscin accumulation in nematodes. Environ Toxicol Phar 31(1): 179-188.




This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

[Submit Article](#)

DOI: 10.32474/JBRS.2020.01.000109

JBRS



Journal of Biosensors & Renewable Sources

Open Access Journal of Biosensors & Renewable sources

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles