



# The Gompertz Length Biased Exponential Distribution and its application to Uncensored Data

Obubu Maxwell<sup>1\*</sup>, Oluwafemi Samuel Oyamakin<sup>2</sup> and Eghwerido Joseph Thomas<sup>3</sup>

<sup>1</sup>Department of Statistics, Nnamdi Azikiwe University, Awka, Nigeria

<sup>2</sup>Department of Statistics, University of Ibadan, Ibadan, Nigeria

<sup>3</sup>Department of Mathematics and Computer Science, Federal University of Petroleum Resources, Effurun, Nigeria

\*Corresponding author: Obubu Maxwell, Department of Statistics, Nnamdi Azikiwe University, Awka, Nigeria

Received: 📅 February 22, 2019

Published: 📅 March 08, 2019

## Abstract

This paper proposes a generalization of the length biased exponential distribution, called the Gompertz length biased exponential (GLBE) distribution. Some of the basic properties of the proposed model were derived in minute details and model parameters estimated by the maximum likelihood estimate method. The adequacy of the model is empirically validated with the use of real - life data.

**Keywords:** Exponential Distribution; Length Biased; Gompertz Generalized Family Of Distribution; Quantile Function; Hazard Functions; Survival Function

## Introduction

Length biased distributions are special case of the more general form known as weighted distribution [1], first introduced by [2] to model ascertainment bias and formalized in a unifying theory by [3]. Lifetime data may be modeled with several existing distributions, although the existing models are not adequate or are less representative of actual data in many situations. Therefore, the development of compound distributions that could better describe certain phenomena and make them more flexible than the baseline distribution is of great importance [4]. Thus, the choice of the model is also an important issue for reliable model parameter estimation. Some exponential distribution generalizations for modeling lifetime data due to some interesting advantages have been recently proposed [5]. In recent years many exponential distribution generalizations have been developed, such as the Marshall Olkin length biased exponential distribution [5], exponentiated exponential [6,7], generalized exponentiated moment exponential [8], extended exponentiated exponential [19], Marshall-Olkin exponential Weibull [10], Marshall-Olkin generalized exponential [5], and exponentiated moment exponential [11] distributions.

A random variable  $X$  is said to have a length biased exponential distribution with parameter  $\beta$  if its probability density function (pdf) and cumulative distribution function (cdf) is given by equation (1) and (2) respectively [12]:

$$g(x, \beta) = \frac{x}{\beta^2} e^{-x/\beta}, x > 0, \beta > 0 \quad (1)$$

$$G(x) = 1 - \left(1 + \frac{x}{\beta}\right) e^{-x/\beta}, x > 0, \beta > 0 \quad (2)$$

Where  $\beta$  is the scale parameter.

The survival function is given by the equation

$$S(x) = \left(1 + \frac{x}{\beta}\right) e^{-x/\beta}, x > 0, \beta > 0 \quad (3)$$

The hazard function is

$$H(x) = \frac{\frac{x}{\beta} e^{-x/\beta}}{\left(1 + \frac{x}{\beta}\right) e^{-x/\beta}}, x > 0, \beta > 0 \quad (4)$$

And the reversed hazard rate function is

$$R(x) = \frac{\frac{x}{\beta^2} e^{-x/\beta}}{1 - (1 + \frac{x}{\beta}) e^{-x/\beta}}, x > 0, \beta > 0 \tag{5}$$

Alzaatreh et al. [13] defined the cumulative distribution function of the Transformed-Transformer (T-X) family of distributions by;

$$F(x) = \int_a^{w[G(x)]} r(t) dt \tag{6}$$

And the corresponding probability density function by;

$$f(x) = \left\{ \frac{d}{dx} w[G(x)] r\{w[G(x)]\} \right\} \tag{7}$$

Morad Alizadeh et al [14] defined the cumulative distribution function and probability density function of the Gompertz Generalized family of distribution by setting  $w[G(x)] = -\log[1 - G(x)]$  and  $r(t) = \theta e^{\gamma t} e^{(\gamma t - 1)}$ ,  $t > 0$  as;

$$F(x) = \int_0^{-\log[1-G(x)]} \theta e^{\gamma t} e^{-(e^{\gamma t} - 1)} dt = 1 - e^{-\frac{\theta}{\gamma} \{1 - G(x)^{-\gamma}\}}, \theta > 0, \gamma > 0 \tag{8}$$

And

$$f(x) = \theta g(x) [1 - G(x)]^{-\gamma - 1} e^{-\frac{\theta}{\gamma} \{1 - G(x)^{-\gamma}\}}, \theta > 0, \gamma > 0 \tag{9}$$

respectively. Where  $\theta$  and  $\gamma$  are additional shape parameters whose role is to vary the tail length.

Thus, we proposed a new generalization of the length biased exponential distribution called the Gompertz length biased exponential (Go-LBE) distribution. In the rest of the paper, we define the Go-LBE model and plots for different parameter values in Section 2; some of the statistical properties of the proposed Go-LBE distribution are discussed in minute details in section 3, Application of the Go-LBE distribution to a lifetime data in section 4. The concluding remark is presented in section 5.

### Gompertz Length Biased Exponential (Go-LBE) Distribution

The cumulative distribution function of the

$$Go-LF_{Go-LBE}(x) = 1 - e^{-\frac{\theta}{\gamma} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\}}, \theta > 0, \gamma > 0, \beta = 0 \tag{10}$$

The corresponding probability density function is given by

$$f_{Go-LBE}(x) = \theta \frac{x}{\beta^2} e^{-x/\beta} \left[ (1 + \frac{x}{\beta}) e^{x/\beta} \right]^{-\gamma - 1} e^{-\frac{\theta}{\gamma} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\}}, \theta > 0, \gamma > 0, \beta = 0 \tag{11}$$

The survival function is given by

$$S_{Go-LBE}(x) = e^{-\frac{\theta}{\gamma} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\}}, \theta > 0, \gamma > 0, \beta = 0 \tag{12}$$

The hazard, reverse hazard and cumulative hazard function is given by equation (13), (14), (15) respectively

$$h_{Go-LBE}(x) = \theta \frac{x}{\beta^2} e^{-x/\beta} \left[ (1 + \frac{x}{\beta}) e^{x/\beta} \right]^{-\gamma - 1}, \theta > 0, \gamma > 0, \beta = 0 \tag{13}$$

$$\gamma_{Go-LBE}(x) = \frac{\theta \frac{x}{\beta^2} e^{-x/\beta} \left[ (1 + \frac{x}{\beta}) e^{x/\beta} \right]^{-\gamma - 1} e^{-\frac{\theta}{\gamma} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\}}}{1 - e^{-\frac{\theta}{\gamma} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\}}}, \theta > 0, \gamma > 0, \beta = 0 \tag{14}$$

$$H_{Go-LBE}(x) = -\log \frac{\theta e^{-\frac{\theta}{\gamma} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\}}}{1 - (1 - \theta) e^{-\frac{\theta}{\gamma} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\}}}, \theta > 0, \gamma > 0, \beta = 0 \tag{15}$$

The odd of the Go-LBE distribution is given by;

$$O_{Go-LBE}(x) = \frac{1 - e^{-\frac{\theta}{\gamma} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\}}}{e^{-\frac{\theta}{\gamma} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\}}}, \theta > 0, \gamma > 0, \beta = 0 \tag{16}$$

The plots for different parameter values of the Go-LBE distribution are given in Figure 1-6 below




Figure 1: Graph for Go-LBE cumulative distribution function at different parameter values.




Figure 4: Graph for Go-LBE hazard function at different parameter values.




Figure 2: Graph for Go-LBE probability density function at different parameter values.




Figure 5: Histogram of the fitted distributions.




Figure 3: Graph for Go-LBE survival function at different parameter values.




Figure 6: Empirical cdf of the fitted distributions.

### Some Statistical Properties of the Go-LBE Distribution

Basic properties such as the asymptotic behavior, parameter estimation and order statistics of the Go-LBE distribution are discussed in minute details.

#### Asymptotic Behavior

Here we critically examine the behavior of the Go-LBE model in equation (11) as  $x \rightarrow 0$  and as  $x \rightarrow \infty$

$$\lim_{x \rightarrow 0} f_{Go-LBE}(x) = \lim_{x \rightarrow 0} \frac{x}{\beta^2} \theta e^{-x/\beta} \left[ \left(1 + \frac{x}{\beta}\right) e^{x/\beta} \right]^{-\gamma-1} e^{\frac{\theta}{\gamma}} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\} = 0$$

$$\lim_{x \rightarrow \infty} f_{Go-LBE}(x) = \lim_{x \rightarrow \infty} \frac{x}{\beta^2} e^{-x/\beta} \left[ \left(1 + \frac{x}{\beta}\right) e^{x/\beta} \right]^{-\gamma-1} e^{\frac{\theta}{\gamma}} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\} = 0$$

This indicates that the Gompertz length biased exponential distribution is unimodal. A clear observation of Figure 2 shows the Go-LBE model has only one peak. This supports our claim that the Go-LBE distribution has only one mode.

#### Parameter Estimation

Using maximum likelihood estimation techniques, we estimate the unknown parameter of the Go-LBE model based on a complete sample. Let  $X_1, \dots, X_n$  indicate a random sample of the complete Go-LBE distribution data, and then the sample's likelihood function is given as;

$$L = \prod_{i=1}^n f(x_i; \theta, \beta, \gamma)$$

$$L = \prod_{i=1}^n \theta \frac{x_i}{\beta^2} e^{-x_i/\beta} \left[ \left(1 + \frac{x_i}{\beta}\right) e^{x_i/\beta} \right]^{-\gamma-1} e^{\frac{\theta}{\gamma}} \{1 - [(1 + \frac{x_i}{\beta}) e^{-x_i/\beta}]^{-\gamma}\}$$

We can now express the log likelihood function as;

$$L(x) = n[\ln \theta - 2 \ln \beta + \frac{\theta}{\gamma}] - \sum_{i=1}^n \left(\frac{x_i}{\beta}\right)^{-\gamma-1} - (\gamma + 1) \sum_{i=1}^n \left(1 + \frac{x_i}{\beta}\right) + \sum_{i=1}^n \ln[1 - \{(1 - \frac{x_i}{\beta}) \exp(\frac{-x_i}{\beta})\}^{-\gamma}] \tag{17}$$

By taking the derivative with respect to  $\theta, \gamma$  and  $\beta$ , and fixing the outcome to zero, we have;

$$\frac{\partial L(x)}{\partial \theta} = n \left[ \frac{1}{\theta} + \frac{1}{\gamma} \right] = 0 \tag{18}$$

$$\frac{\partial L(x)}{\partial \gamma} = \frac{-n\theta}{\gamma^2} - \sum_{i=1}^n \left(\frac{x_i}{\beta}\right)^{-\gamma} \ln\left(\frac{x_i}{\beta}\right)^{-\gamma} - \sum_{i=1}^n \left(1 + \frac{x_i}{\beta}\right) - \frac{\sum_{i=1}^n \{(1 + \frac{x_i}{\beta}) \exp(\frac{-x_i}{\beta})\} \{(\frac{-x_i}{\beta}) \ln(\frac{x_i}{\beta})\}}{1 - \{(1 + \frac{x_i}{\beta}) \exp(\frac{-x_i}{\beta})\}^{-\gamma}} = 0 \tag{19}$$

$$\frac{\partial L(x)}{\partial \beta} = \frac{-2n}{\beta} + \sum_{i=1}^n \left(\frac{x_i}{\beta^2}\right) + (\gamma + 1) \sum_{i=1}^n \left(\frac{x_i}{\beta^2}\right)^{-(\gamma+2)} \left(\frac{x_i}{\beta^2}\right) - \frac{\sum_{i=1}^n \{(1 + \frac{x_i}{\beta}) \exp(\frac{-x_i}{\beta})\}^{-(\gamma+1)} \{(\frac{-x_i}{\beta}) \exp(\frac{x_i}{\beta})\}}{1 - \{(1 + \frac{x_i}{\beta}) \exp(\frac{-x_i}{\beta})\}^{-\gamma}} \tag{20}$$

Solving equation (18)-(20) iteratively, will give the estimate of the parameters of the Go-LBE model.

#### Order Statistics

We considered a random sample denoted by from the densities of the Go-LBE distribution. Then,

$$f_{j:n}(x) = \frac{n!}{(j-1)!(n-1)!} f_{Go-LBE}(x) f_{Go-LBE}(x)^{j-1} [1 - F_{Go-LBE}(x)]^{n-j}$$

The probability density function of the order statistics for the Go-LBE distribution is given as;

$$f_{j:n}(x) = \frac{n!}{(j-1)!(n-j)!} \theta \frac{x}{\beta^2} e^{-x/\beta} \left[ \left(1 + \frac{x}{\beta}\right) e^{x/\beta} \right]^{-\gamma-1} e^{\frac{\theta}{\gamma}} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\} \left[ 1 - e^{\frac{\theta}{\gamma}} \{1 - [(1 + \frac{x}{\beta}) e^{-x/\beta}]^{-\gamma}\} \right]^{n-j} \tag{21}$$

The Go-LBE distribution has minimum order statistics given as;

$$f_{1:n}(x) = n\theta \frac{x}{\beta^2} e^{-x/\beta} \left[ (1 + \frac{x}{\beta}) e^{-x/\beta} \right]^{\gamma-1} e^{-x/\beta} \left[ (1 + \frac{x}{\beta}) e^{-x/\beta} \right]^{\gamma} e^{-x/\beta} \left[ (1 + \frac{x}{\beta}) e^{-x/\beta} \right]^{-n-1} \quad (22)$$

And maximum order statistics given as;

$$f_{n:n}(x) = n\theta \frac{x}{\beta^2} e^{-x/\beta} \left[ (1 + \frac{x}{\beta}) e^{-x/\beta} \right]^{\gamma-1} e^{-\theta/\gamma} \left[ (1 + \frac{x}{\beta}) e^{-x/\beta} \right]^{\gamma} 1 - e^{-\theta/\gamma} \left[ (1 + \frac{x}{\beta}) e^{-x/\beta} \right]^{-n-1} \quad (23)$$

## Data Analysis

Here, we provide an application of the Gompertz length biased exponential distribution by comparing the results of the model fit with that of other Gompertz- G family of distributions. The data set we employ is the uncensored strength of 1.5cm glass fibre data

previously used by Bourguignon M et al. [15], Merovci F et al. [16]. This data set will be used to compare between fits of the Gompertz length biased exponential distribution (Go-LBE) with that of Gompertz-Exponential (Go-E), Gompertz-Lomax (Go-L), and, Gompertz-Weibull (Go-W). The data is presented below (Tables 1 & 2):

**Table 1:** Descriptive Statistics on Cancer Stem Cell Data.

Min.	1 <sup>st</sup> Qu.	Median	Mean	3 <sup>rd</sup> Qu.	Max.	Skewness	Kurtosis	Variance
0.550	1.375	1.590	1.507	1.685	2.240	-0.8999263	3.923761	0.1050575

**Table 2:** MLEs, SW, AD and K-S of parameters for Cancer Stem Cell data.

Model	Estimates				SW	K-S	AD
	$\beta^*$	$\theta^*$	$\gamma^*$	$\delta^*$			
Go-LBE (Proposed Model)	0.125465239	5.951201245	1.251202365	-	0.1209	0.1195	0.7845
Go-E	-0.004768848	-1.810999364	-1.987714978	-	0.1445	0.1314	0.8425
Go-L	0.004592168	8.179090955	0.506999370	1.515829085	0.1685	0.1542	0.9462
Go-W	0.228488761	0.009628097	0.794918813	5.612111282	0.2329	0.1520	1.2832

0.55, 0.74, 0.77, 0.81, 0.84, 1.24, 0.93, 1.04, 1.11, 1.13, 1.30, 1.25, 1.27, 1.28, 1.29, 1.48, 1.36, 1.39, 1.42, 1.48, 1.51, 1.49, 1.49, 1.50, 1.50, 1.55, 1.52, 1.53, 1.54, 1.55, 1.61, 1.58, 1.59, 1.60, 1.61, 1.63, 1.61, 1.61, 1.62, 1.62, 1.67, 1.64, 1.66, 1.66, 1.66, 1.70, 1.68, 1.68, 1.69, 1.70, 1.78, 1.73, 1.76, 1.76, 1.77, 1.89, 1.81, 1.82, 1.84, 1.84, 2.00, 2.01, 2.24

data set, Table 2 shows parameter estimate and the value for the Shapiro Wilk (S-W), Anderson Darling (AD), and the Kolmogorov Smirnov (K-S) statistic (Table 3).

From Table 3, the Go-LBE has the highest log-likelihood values and the lowest AIC, CAIC, BIC and HQIC values; hence, it is chosen as the most appropriate model amongst the considered distributions.

For all competing distributions using the strength of glass fibre

**Table 3:** Log-likelihood, AIC, AICC, BIC and HQIC values of models fitted for Cancer Stem Cell data.

Model	Negative LL	AIC	CAIC	BIC	HQIC	P-VALUE
Go-LBE (Proposed Model)	14.01129	32.5215	34.5941	41.1105	37.1523	0.3542
Go-E	14.81765	35.6353	36.04208	42.06471	38.16402	0.227104
Go-L	14.50274	37.00548	37.69513	45.57802	40.3771	0.099874
Go-W	15.18847	38.37694	39.06659	46.94948	41.74856	0.108882

## Conclusion

This research has successfully extended the length biased exponential distribution. Densities and basic statistical expressions were briefly derived. The performance of the proposed Gompertz length biased exponential distribution was compared to existing models in literature based on the negative log likelihood, AIC, CAIC, BIC and HQIC values. Based on the lowest criterion values, we therefore conclude that the Gompertz length biased exponential

distribution is the most suitable model amongst the considered models and indeed a very competent model for describing life-time situations.

## References

1. Kareema Abed Al-kadim, Nabeel Ali Hussein (2014) New proposed length biased weighted exponential and reyleigh distribution with applications. Mathematical theory and Modeling 4(7): 137-152.
2. RA Fisher (1934) The effects of methods of ascertainment upon the estimation of frequency. Ann. Eugen 6(1): 13-25.

3. CR Rao (1965) On discrete distributions arising out of methods of ascertainment. In *Classic. Contag Discrete Distributions*. Statistical G.P Patil, Edn, Publishing Society, Calcutta, India pp. 320-332.
4. Maxwell O, Friday AI, Chukwudike NC, et al. (2019) A theoretical analysis of the odd generalized exponentiated inverse Lomax distribution. *Biom Biostat Int J*. 8(1): 17-22.
5. Ristic MM, Kundu D (2015) Marshall-Olkin generalized exponential distribution, *Metron* 73(3): 317-333.
6. Gupta RD, Kundu D (1999) Theory & methods: Generalized exponential distributions. *Australian & New Zealand J. Stat* 41 (2): 173-188.
7. Gupta RD, Kundu D (2001) Exponentiated exponential family: An alternative to gamma and Weibull distributions. *Biometrical J* 43 (1): 117-130
8. Iqbal Z, Hasnain SA, Salman M, Ahmad M, Hamedani G (2014) Generalized exponentiated moment exponential distribution. *Pakistan J. Stat* 30(4): 537-554.
9. Abu-Youssef S, Mohammed B, Sief M (2015) An extended exponentiated exponential distribution and its properties. *Int. J. Comput. Appl* 121(5): 1-6
10. Pogány TK, Saboor A, Provost S (2015) The Marshall-Olkin exponential Weibull distribution. *Hacetatepe J. Math. Stat* 44(6): 1579-1594
11. Hasnain SA, Iqbal Z, Ahmad M (2015) On exponentiated moment exponential distribution. *Pak. J. Statist.* 31(2): 267-280.
12. Dara ST, Ahmad M (2012) Recent Advances in Moment Distribution and Their Hazard Rates. *Lap Lambert Academic Publishing GmbH KG* pp. 128.
13. Alzaatreh A, C Lee, F Famoye (2013) A new method for generating families of continuous distributions. *Metron* 71 (1): 63-79.
14. Morad Alizadeh, Gauss M Cordeiro, Luis Gustavo Bastos Pinho, Indranil Ghosh (2017) The Gompertz-G family of distributions, *Journal of Statistical Theory and Practice* 11(1): 179-207.
15. Bourguignon M, Silva RB, Cordeiro GM (2014) The Weibull-G family of probability distributions. *Journal of Data Science* 12: 53-68.
16. Merovci F, Khaleel MA, Ibrahim NA, Shitan M (2016) The beta type X distribution: Properties with applications. *Springer Plus* 5: 697.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

[Submit Article](#)

DOI: 10.32474/CTBB.2019.01.000111



### Current Trends on Biostatistics & Biometrics

#### Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles