



Anti-Microbial and Antioxidant Activities of Ziziphus spina Christi, Sudanese Medicinal Plant

Mahdi Abdu-Almageed Mohamed Ali¹, Asaad Alsiddig Ahmed², Ali Abdurrahman Almakki³ and Hatim MY Hamadnalla^{4*}

¹Department of Biology and Technology, College of Applied Sciences - University of Bahri, Sudan

²Department of Biology - Khartoum International Preparatory School, Sudan

³Department of Microbiology, College of Applied Sciences- University of Bahri, Sudan

⁴Department of Biochemistry, College of Applied Sciences, University of Bahri, Sudan

*Corresponding author: Hatim MY Hamadnalla, Department of Biochemistry, College of Applied Sciences, University of Bahri, Sudan

Received: 📅 April 28, 2020

Published: 📅 June 25, 2020

Abstract

This work is an anti-Microbial and Antioxidant activity of Ziziphus spina Christi based on four selected bacteria: Escherichia coli, Pseudomonas aeruginosa, Bacillus subtilis and Staphylococcus aureus. These species were collected and prepared in Sudan. The concentrations were carried out and compared with standard control sample, the suspension of anti-microbial activities have shown different results in Fagonia cretica. The antioxidant activity is the ability of bioactive compounds to prevent, delay and protect against oxidant of virus substrates such as DNA and lipid materials, both in living organisms (e. g. human) and in food products. The result showed an antioxidant activity based on standard liquid reagent [1].

Introduction

Anti- Microbial is an agent that kills microorganisms such as bacterial or stops their growth. Anti-microbial medicines can be grouped according to the microorganisms they act primarily against. For example: antibiotics were used against bacteria and antifungals too [2]. Anti- Microbial can be further subdivided into bactericidal agent; instead, certain types of porous media have been developed to kill microbes on contact [3]. On the other hand, antioxidants are compounds that inhibit oxidation. This is a chemical reaction that can produce free radicals. Thereby, leading to chain reactions that may damage the cells of organisms. For example: thiols or ascorbic acid terminates these chain reactions. Mostly, antioxidants have two entirely different groups of substances: industrial chemical reactions which can be added to products and natural chemicals which can be presented in foods and tissues for resistant of cancer [4]. Plant and animals maintain complex system of overlapping antioxidant [5].

Study area

The batch was found in west Om-Dorman region, Sudan. This area is mostly populated and features a hot arid climate routinely

exceeded to 40 °C in midsummer, the region lay between 15.6476° N, 32.4807°E. This region considered as heliostat city of Sudan. The soil of this area is partly divided into two types: slab stone and muddy.

Population

According to ethno-botanical Omdurman area includes several people such as Arabs and Africans. Meanwhile, most of these people are officers beside other traditional works such as agriculture and animal husbandry.

Methodology

Experimental materials

Plant material was collected from west Om-Dorman region, Khartoum - Sudan in January 2020. The identification of the plant was carried out at natural research center and compared with herbarium of Department of Biology and Technology - Faculty of applied Science and industry - University of Bahri, Sudan. The medicinal plant was dried under dark shade for one week.

Preparation of crude extract

One hundred gm. of whole plant material (Sedir - local name) was extracted by ethyl acetate for (24) hours at room temperature. The extracts were filtered using filter paper and air-dried. The final volume was adjusted to give the specific concentration. Yield of all extract was respected to dry starting materials.

Anti-microbial activity (preparation of culture)

Twenty-eight gm. of nutrient agar media was added to (1000) ml. of distilled water. Seven gm. was added to (250) ml. distilled water on glass bottle volume (250) ml. (4 bottles), the four bottles were shaken very well to dilute the powder media and (32.5) gm. of sabouraud dextrose agar was added to (500) ml. distilled water. Meanwhile, (16.25) gm. was added to (250) gm. distilled water on glass bottle volume (250) ml. then all bottles were shaken and put on autoclave for sterilization at (121°C) and (15 lbs) pressure for (15) minutes.

Preparation of sabouraud

Sixty two gm. of powdered sabouraud dextrose agar was weighted and dispersed in (1) liter of distilled water and allowed to soak for (10) minutes swirl to mixed and sterilized by autoclave for (15) minutes at (121°C) and cooled to (47 °C). Lastly, mixed well and poured into petri dishes.

Bacteria organisms test

- a) Staphylococcus aureus (ATCC 6538 Gram +ve Bacteria).
- b) Bacillus subtilis (ATCC 6633 Gram +ve Bacteria).
- c) Pseudomonas aeruginosa (ATCC 8739 Gram -ve Bacteria).
- d) Escherichia coli (ATCC 8739 Gram -ve Bacteria).

Preparation of organism's suspension

One ml. aliquots of a (24) hours broth culture of the testing organisms were aseptically distributed on nutrient agar slopes and incubated at (37 °C) for (24) hours, the bacteria growth was then harvested and washed off with (100) ml. sterile normal saline to produce a suspension containing about (108 - 109) C.F.U/ml., the suspension was then stored in a refrigerator at (4 °C). The average number of organisms per ml. of the stock suspension was determined by the surface counting technique. (Miles and Misra, 1938). Serial dilutions of the stock suspension were made in sterile normal saline solution and (0.02) ml. volumes of the appropriate dilutions were transferred by micro-pipette onto the surface of dried nutrient agar plates, the plates were allowed to stand for (2) hours at room temperature with drops to dry and incubated at (37 °C) for (2) hours, after incubation the number of developed colonies in each drop was counted. The average number of the colonies per drop of (0.02) ml. was multiplied by (50) and by dilution factor to give viable count of the stock suspension which expressed as a number of colony of the stock suspension which expressed as a

number of colony forming unit (C.F.U). Therefore, each time of fresh stock suspension was prepared.

Testing for anti-microbial activity

The cup-plate agar diffusion method was adopted with some minor modifications to assess the anti-bacterial activity of the prepared extract [6]. One ml. of standardized bacterial stock suspension 108 - 109 C.F.U/ml was thoroughly mixed with (100) ml. of molten sterile nutrient agar which was maintained at (45 °C). Twenty ml. aliquots of inoculated nutrient agar were distributed into sterile petri-dishes, the agar was left to set in each of this four cup-plates (10 mm. in diameter) were cut using a sterile cork borer and agar was removed. Alternate cups were filtered with (0.1) ml. sample of each of the oil dilutions in methanol using automatic micro-liter pipette at room temperature for (2) hours. The plates were then incubated in the upright position at (37 °C) for (18) hours. Two replicates were carried out for each extract against each of the test organisms, after incubation the diameter of the resultant growth inhibition zones were measured, averaged and mean values were tabulated.

DPPH radical scavenging assay

The DPPH assay scavenging was determined according to the method of Shimada [1] with some modification. In 96-well plate, the test samples were allowed to react with 2,2-Di (4-tert-octyl phenyl)-1-picryl-hydrazyl stable free radical (DPPH) for half an hour at 37 °C. The concentration of DPPH was kept as (300µM). The test samples were dissolved in DMSO while DPPH was prepared in ethanol. After incubation, decrease in absorbance was measured at (517nm) using multiplate reader spectrophotometer. Percentage radical scavenging activity by samples was determined in comparison with a DMSO treated control group. All tests and analysis were run in triplicate.


Discussion

Anti-Microbial

Ethyl acetate solvent had been detected a precise output. The numbers of bacteria were infected by concentrations Table 1. The dead bacteria were increased when we use anti-biotic. On other hand, *P. aeruginosa* and *B. subtilis* resulted the same dead bacteria against anti-biotic although they were different gram stains, while *E. coli* bacteria resulted the lowest number of dead bacteria. The (Table 1) also shows the increasing of concentrations have led to more infected bacteria. These results had been depended on the cell wall strengthen of the bacteria. The standard control samples had mentioned the numbers of colonies and the teared bacteria which were measured by (1) gm /ml. concentration. Several bioassay such as disk diffusion, well diffusion and broth or agar dilution are well known and commonly used, but others such as flow cytometric and bioluminescent methods are not widely use because they requires specified equipment, evaluation for reproducibility and standardization [7].

Table 1: Anti-microbial activities of Ziziphus spina Christi plant against standard bacterial strains at concentration 1 gm/ml.

Plant extract	Concentrations	Standard bacterial strains			
		<i>E. coli</i> (-ve)	<i>Paeruginosa</i> (-ve)	<i>B. subtilis</i> (+ve)	<i>S.aureus</i> (+ve)
Ethyl acetate	100	16	19	22	21
	50	15	18	19	20
	25	14	17	19	19
	12.5	12	15	17	17
	amoxicillin	24	36	36	34



Antioxidant

Propyl gallate was used as standard and showed the result of (31±0.03) in Table 2 after running of triplicate process. That means Ziziphus spina Christi has an antioxidant activity through

the comparison with dimethyl sulfoxide (DMSO) which considered as oregano sulfur compound. This colorless liquid is an important polar aprotic solvent that dissolves both polar and non-polar compounds and is miscible in wide range of organic solvents as well as water. It has a relatively high boiling point [8].

Table 2: The antioxidant activity of Ziziphus spina Christi extract against standard propyl gallate.

No.	% RSA ± SD (DPPH)
Z.S ethylacetate Extract	61 ± 0.04
Standard (Propyl gallate)	93 ± 0.01

Also, propyl gallate exhibits antimicrobial activity and has reported to an effective antioxidant based hepatoprotector. Finally, this reagent used as anti-fade in fluorescence microscopy to reduce photobleaching of fluorescent probes such as rhodamine and fluorescein [1]. A balance between free radical and antioxidant is necessary for proper physiological function. If free radical overwhelms the body's ability to regulate them a condition known as oxidative stress ensues. Application of external source of antioxidants can assist in coping oxidative stress [9-11].

Conclusion

Anti-microbial

- Anti-biotics had the ability to inhibit the growth of bacteria; therefore, we must detect the suitable one for treatment Table 1.
- Anti-bacterial activity differs with applied extractive method.

- Anti-bacterial activities support further studies to discover the new chemical structures that can contribute to alleviate or cure some illnesses.

Antioxidant

- Antioxidants counteract by products called free radical.
- Antioxidant may possible the development of cancer.
- Some food sources of antioxidant such as Beta-Carotene, vitamin (A), (C) and (E) must be available within the meals.

Acknowledgement

- Great thanks for natural research center for their cooperation.
- Deep greetings to the Department of Biology and Technology - Faculty of Applied science and Technology and Industry - University of Bahri.

References

1. Shimada K, Fujikawa K, Yahara K, Nakamura T (1992) Antioxidative properties of xanthan on the anti-oxidation of soya bean oil in cyclodextrin emulsion 40: 945-948.
2. Wain W (1989) Mould in ancient and more recent medicine 3(1): 21-23.
3. Kingston W (2008) Irish contributions to the origins of antibiotics 177(2): 87-92.
4. Jiang L, Pearson, Crews ST (2010) Diverse mode of *Drosophila* tracheal fusion cell transcriptional regulation 127(5-6): 265-280.
5. Stanner SA, Hughes J, Kelly CNM, J Buttriss (2004) A review of epidemiological evidence for the antioxidant hypothesis 7(3): 22-407.
6. Sabath MD (1976) The assay of antimicrobial compounds 7(3): 287-295.
7. Mounyr B, Moulay S, Saad KI (2016) Methods for in vitro evaluating antimicrobial activity 6(2): 71-79.
8. Smith ER, Hadidian Z, Mason MM (1967) Dose toxicity of dimethyl sulfoxide 141(1): 96-109.
9. Lobo V, Patel A, Phalak A, Chandra N (2010) Free radicals, antioxidants, and functional foods: Impact in human health 4(8): 118-126.
10. Miles AA, Misra SS (1938) The estimation of bactericidal power of the blood 38(6): 732-749.
11. Fleming A (1929) On the antibacterial reaction of cultures of a penicillium with special reference to their use in isolation of *B. influenza* 10(3): 226-236.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

[Submit Article](#)

DOI: [10.32474/OAJESS.2020.05.000204](https://doi.org/10.32474/OAJESS.2020.05.000204)



Open Access Journal of Environmental and Soil Sciences

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles