



A Review on Physical and Functional Properties of *Lepidium sativum* L

Manju* and Neetu Dobhal

Department of Foods and Nutrition, College of Home Science, Govind Ballabh Pant University of Agriculture and Technology, India

*Corresponding author: Manju, Department of Foods and Nutrition, College of Home Science, Govind Ballabh Pant University of Agriculture and Technology, India

Received:  June 10, 2022

Published:  June 20, 2022

Introduction

Garden cress is an annual herb that grows in the Middle East, Europe, and the United States. It belongs to the Brassicaceae family and the Kingdom Plantae. Garden cress is thought to have originated in Ethiopia, according to scientific evidence. Garden cress can thrive in any climate and soil type. The seeds resemble some oil seeds in appearance, with dicotyledonous endosperm accounting for 80-85 percent of the seed content and the seed coat and embryo accounting for 12-17 percent and 2-3 percent, respectively [1,2]. Garden cress seeds are used in a variety of foods as a gelling agent, a water retention agent, a thickening agent, an emulsifying agent, and a binding agent. As a result, the purpose of this review is to compile the physical and functional properties of garden cress seeds. The present article is mainly divided into two sections

- a) Physical Characteristics of Garden Cress Seeds
- b) Functional Characteristics of Garden Cress Seeds Flour

Physical Characteristics of Garden Cress Seeds

Garden cress seeds are small oval-shaped, pointed and triangular at one end, smooth, about 2-3 mm long, 1-1.5 mm wide, slightly broader at base and notched at apex, dorsiventrally bulging, with a furrow present on both surfaces extending up to two thirds downward, and a slight wing-like extension present on both edges of seed. By absorbing water, the seeds swell and become slippery when combined with water, and after boiling, they form a gel-like material with strong adhesive properties [3] and [4]. According to Paranjape and Mehta [5] garden cress seeds are morphologically similar to oilseeds, with the dicotyledonous endosperm accounting for 80-85 per cent of the seed matter, the seed coat 12-17 percent, and the embryo 2-3 percent. Gokavi et al. [6] conducted study on physical composition of garden cress seeds and found that the seeds were brownish red in colour and oval in shape. The seed length and width were $298 \pm 3.2 \mu\text{m}$ and $100 \pm 1.9 \mu\text{m}$ respectively. The weight, volume and bulk density of one thousand kernels was $2.5 \pm 0.13\text{g}$, $3.3 \pm 0.15 \text{ml}$ and $0.75 \pm 0.01\text{g/ml}$, respectively.

A study of Vaishnavi et al. [7] revealed that garden cress seeds were oval shaped and reddish-brown in colour. Dimensional properties viz. length, width and thickness measured using vernier calliper were found as 2.6mm, 1.2mm and 0.94mm, respectively. Sphericity was 54.59 per cent and the average weight of 1000 garden cress seeds was 1.86g. True density and bulk density of garden cress seeds were measured using a graduated cylinder and were calculated as 1230 kg/m^3 and 729.74 kg/m^3 respectively. Porosity of garden cress seeds was 40.67% and angle of repose for garden cress seeds was calculated as 20.590.

Patel et al. [8] studied the physicochemical properties of *Lepidium sativum* mucilage powder in which they found that particle size (μm), swelling ratio, viscosity, bulk density (g/ml) and tapped density (gm/ml) were 189.57, 3.7, 8.05, 0.37 and 0.62 respectively. Najia et al. [9] analyzed the influence of thermal treatments on textural characteristics of cress seed gum gel. In this study, the cress seeds were treated with different thermal treatments viz. 600C for 30min, 800C for 23min, 1000C for 18min and 1210C for 15min. Textural stability were studied by using penetration and Instrumental Texture Profile Analysis (TPA) tests. The obtained results revealed that cress seed gel was heat stable in almost all of the textural attributes and gel at 7 per cent concentration had a cohesiveness value of 0.43, hardness 11 g, consistency 77 g, adhesiveness 9.2 g, elasticity 11.23 g/s, springiness 3.8 mm, gumminess 44 g, chewiness 170 gmm. Wadhwa et al. [10] found that garden cress seeds were ovoid and flattened, measuring 2-3 mm in length and varied in colour from pale brown to brownish red to black. The seedling germinated epigeally; the cotyledons were tri-foliolate, the leaflets were spatulate, and the lateral leaflets were smaller than the central one. Deshmukh et al. [11] observed true density and bulk density of garden cress seeds as 0.76 and 1.25 g/ml, respectively. The porosity of seed was 39.2 per cent and angle of repose was 190. The bulk and tapped densities of dried garden cress seeds mucilage powder before and after roasting process was examined by Hassan and Abdel-Rahman [12]. The bulk density of mucilage extracted from raw and roasted

sample was recorded as 0.1396 and 0.17013 g/ml, respectively. The tapped density was 0.2178 and 0.3162 g/ml, respectively for raw and roasted samples. They found that increase in bulk and tapped density was directly proportional to roasting time. Sonawane et al. [13] studied the physicochemical characteristics of extracted cress seed mucilage and reported the swelling index, bulk density, tapped density, compressibility index, pH, extractive values, particle length and breadth as 11ml, 0.2857 g/cc, 0.3389 g/cc, 15.69 per cent, 5.6, 13.36 per cent, 1.3674 mm and 0.875 mm, respectively. Gaikwad et al. [14] reported 1000 kernel weight, true density, bulk density, porosity and angle of repose of garden cress seeds as 1.96 g, 1182 kg/ m³, 0.76 g/ml, 36 per cent and 25.170 respectively. Different dimensional properties like length were 2.6, width was 1.2 and thickness was measured 0.94 (mm).

Functional Properties of Garden Cress Seeds Flour

Functional properties come under the essential physicochemical properties of foods that explain the interactions between structures, molecular conformation, compositions, and physicochemical properties of food components with the nature of the environment and the conditions under which these are examined. Functional properties are needed to effectively determine and precisely evaluate how different proteins, fats, carbohydrates (starch and sugars), and fibres may function in specific food systems [15,16]. Toliba and Mohamed [17] studied the functional properties i.e., water holding capacity, oil holding capacity, emulsifying activity and stability of garden cress seeds powder and reported these as 4.51±0.83 ml/g, 2.79±0.32 ml/g, 44.54±1.35 per cent and 40.65±1.02 per cent, respectively. Hanan et al. [18] reported the water absorption capacity and oil absorption capacity of garden cress seeds flour as 11.21 g/g and 0.726 g/g, respectively. Kilor and Bramhe [19] reported the water retention capacity of garden cress seeds as 10.5ml/g found with precipitation of soaked and blended seed in acetone and 16.3ml/g in alcohol. Emulsifying and foaming properties of cress seeds gum (1% w/w) were measured by Najia et al. [9]. The observed results revealed that garden cress seeds gum solution has high emulsifying (92 percent) and foaming (25.4 percent) capacity at 250C. Functional properties of protein isolated from garden cress seeds were studied by Ali [20]. The water absorption, fat absorption, emulsification capacity and forming capacity of garden cress seeds were 229±2.89, 350±0.99, 65±3.02 and 39.50±1.57 per cent, respectively.

Ghada et al. [21] studied the functional properties and antihyperlipidemic activities of garden cress seed oil. They reported the refractive index of oil as 1.475, peroxide value (meq. peroxide/kg oil) as 2.06, saponification value (KOH/g oil) as 179.5, iodine value (g of I₂ absorbed/ 100g oil) as 120.6 and oxidative stability (hrs) as 30. Zia-Ul-Haq et al. [22] reported the refractive index, specific gravity, unsaponifiable matter, acid value and saponification values of garden cress seeds as 1.47, 0.82, 0.57 µg/g, 1.04 mg KOH/g and 179.03 mg KOH/g, respectively. Rajshri and Haripriya [23] evaluated the effect of roasting on functional properties of garden cress seeds. They found out that water absorption and oil absorption of untreated garden cress seeds powder was 9.56±0.40ml/g

and 7±0.72ml/g respectively, while it was 8.93±0.30ml/g and 7.4±1.07ml/g respectively for roasted garden cress seeds. There was a significant decrease in water absorption (6.58 per cent) and oil absorption (5.71per cent) on dry roasting. The untreated and roasted seed's powder swelling capacity was found as 11.2±0.68 and 12.1±0.69 per cent, respectively. On dry roasting a significant increase (8.0 percent) in swelling capacity was observed. The Bulk density of untreated and roasted seed powder was reported as 1±0.5g/ml and 0.63±0.01g/ ml respectively. 37 percent decrease in bulk density on dry roasting was recorded.

Conclusion

Food physical and functional properties are important in explaining cooking and processing characteristics because they explain how foods perform in a system as a chemical intermediate or as a significant participant of product attributes. Information on the physical and functional characteristics of garden cress seeds provides the data needed to evaluate and maintain the quality of developed products, as well as an indication of how they would behave in the food system. All of the reviews came to the same conclusion: garden cress seeds have good physical and functional properties and can be used in the food industry as a healthy substitute.

References

- Mathews S, Singhal RS, Kulkarni PR (1993) Some physicochemical properties of *Lepidium sativum* (Haliv) seeds. Die Nahrung 37(1): 69-71.
- Gopalan C, Sastri BV, Balasubramanian SC (2000) Nutritive Value of Indian foods. Hyderabad, India: National Institute of Nutrition, Indian Council of Medical Research.
- Doke S, Guha M (2014) Garden cress (*Lepidium sativum* L.) seeds - An important medicinal source. J Nat Prod Plant Resour 4(1): 69-80.
- Sarkar S, Datta S, Ghosh I (2014) Experimental studies on nutritional medicinal role of garden cress seed on animal and human. Int J Med Chem and Anal 4(1): 41-45.
- Paranjape AN, Mehta AA (2006) A study on clinical efficacy of *Lepidium sativum* seeds in treatment of bronchial asthma. Iranian J Pharmacol Ther 5: 55-59.
- Gokavi SS, Malleshi NG, Guo M (2004) Chemical composition of garden cress (*Lepidium sativum*) seeds and its fractions and use of bran as a functional ingredient. Plant Foods Hum Nutr 59(3): 105-111.
- Vaishnavi, Gupta R, Choudhary P (2020) Botanical description of garden cress (*Lepidium sativum* L.) plant and physical characteristics of its seeds. International Journal of Pharmacognosy and Phytochemical Research 9(5): 2424-2428.
- Patel U, Kulkarni M, Undale V, Bhosale A (2009) Evaluation of Diuretic Activity of Aqueous and Methanol Extracts of *Lepidium sativum* Garden Cress (*Cruciferae*) in Rats. Trop J Pharm Res 8(3): 215.
- Najia, S, Razavia SMA, Karazhiyanb H, Koochekia A (2012) Influence of thermal treatments on textural characteristics of cress seed (*Lepidium sativum*) gum gel. EJEAF Che 11(3): 222-237.


10. Wadhwa S, Panwar MS, Agrawal A, Saini N, Patidar LN (2012) A review on *pharmacognostical* study of *Lepidium sativum*. *Advance research in pharmaceuticals and biological* 2(4): 316-323.
11. Deshmukh YR, Thorat SS, Mhalaskar SR (2017) Influence of garden cress seed (*Lepidium sativum L.*) bran on quality characteristics of cookies. *Int J Curr Microbiol Appl Sci* 6(9): 586-593.
12. Hassan MAM, Abdel Rahman AM (2019) The influence of dry roasting process on chemical and nutritional properties of garden cress seeds flour. *Int Adv Res J Sci Eng Technol* 6(11): 73-82.
13. Sonawane MS, Patil AP, Sonawane RO, Pradyumna PI, Patil PG (2019) *Lepidium sativum* characteristics and as a multifaceted polymer: an overview. *Indo Am J Pharm Sci* 6(5): 9470-9480.
14. Gaikwad VG, Chavan UD, Godase SN, Kotecha PM (2021) Studies on nutritional quality of garden cress seed cookies. *Int J Chem Stud* 9(1): 1603-1609.
15. Kaur M, Singh N (2006) Relationships between selected properties of seeds, flours, and starches from different chickpea cultivars. *Int J Food Prop* 9(4): 597-608.
16. Siddiq M, Nasir M, Ravi R, Dolan KD, Butt MS (2009) Effect of defatted maize germ addition on the functional and textural properties of wheat flour. *Int J Food Prop* 12(4): 860-870.
17. Toliba AO, Mohamed A S (2019) The Effect of Garden Cress Seeds Addition on Rheological Properties of Wheat Flour and Chocolate Flavoured Cupcake. *Egyptian Journal of Food Science* 47(2): 187-199.
18. Hanan MAA, Nahla SZ, Abdelaleem MA (2019) Utilization of garden cress seeds (*Lepidium sativum L.*) as natural source of protein and dietary fiber in noodles. *Int J Pharm Res Allied Sci* 8(3): 17-28.
19. Kilor V, Bramhe NN (2014) Development of effective extraction method for *Lepidium sativum* seed mucilage with higher yield. *J Adv Pharm Educ Res* 4(3): 354-360.
20. Ali M (2013) Preparation and characterization of protein isolate and biodiesel from garden cress seed. *European J Chem* 4(2): 85-91.
21. Ghada MY, Heba EG, Abd El S (2014) Study the physico-chemical properties and antihyperlipidemic activities of garden cress seed oil. *Am J Sci* 10(12): 324-330.
22. Zia Ul Haq M, Ahmad S, Calani L, Mazzeo T, Del Rio D, et al. (2012) Compositional study and antioxidant potential of *Ipomoea hederacea* Jacq. and *Lepidium sativum L.* seeds. *Molecules* 17(9): 10306-10321.
23. Rajshri VS, HariPriya A (2018) Effect of processing on selected nutrient profile of garden cress seeds and development of garden cress seed-based muffin. *International journal of academic research and development* 2(3): 1542-1547.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)

DOI: [10.32474/SJFN.2022.04.000188](https://doi.org/10.32474/SJFN.2022.04.000188)



Scholarly Journal of Food and Nutrition

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles