



# Sinus Mycetoma in A Child

C Ndiaye<sup>1\*</sup>, MS Diouf<sup>2</sup>, NF Thiam<sup>2</sup>, A Dieye<sup>1</sup>, N Ndour<sup>1</sup>, A Tall<sup>1</sup>

<sup>1</sup>Department of Otolaryngology-Head and Neck Surgery, Fann Teaching Hospital, Dakar, Senegal

<sup>2</sup>Department of Otolaryngology-Head and Neck Surgery, Idrissa Pouye General Hospital, Dakar, Senegal

\*Corresponding Author: Dr Ciré Ndiaye, Professor of Oto-Rhino-Laryngology, ENT department of FANN hospital, Department of Otolaryngology-Head and Neck Surgery, Fann Teaching Hospital, Dakar, Senegal.

Received: 📅 January 02, 2023

Published: 📅 April 29, 2024

## Abstract

We report the case of a 14-year-old child who presented a unilateral nasal obstruction associated with purulent rhinorrhea. A CT scan of the sinuses showed hyperdense formations filling the sinuses. Endoscopic surgery was used to remove the fungal balls like a tumor.

## Observation

A 14-year-old patient with no particular pathological history consulted for a left nasal obstruction that had been progressively worsening for 2 years. The patient also complained of left purulent rhinorrhea associated with homolateral epistaxis.

### ENT examination revealed:

1. a friable endonasal mass obstructing the left nasal cavity.
2. homolateral purulent rhinorrhea.
3. left suborbital oedema.


The lymph nodes were free. A CT scan of the sinuses revealed a mucoid filling with macrocalcifications in the left maxillary and sphenoidal sinuses (Figure 1). Immune response was normal. Biopsy of the endonasal mass showed an inflammatory polyp. The patient underwent endoscopic surgery. Endoscopy of the left nasal cavity revealed polyps which, after removal, revealed an accumulation and dense conglomeration of fungal hyphae in the left maxillary, ethmoidal and sphenoidal sinuses (Figure 2). The maxillary, ethmoidal and sphenoidal sinuses were opened wide. The post-op

erative period was uneventful. After 4 months' follow-up, there was no recurrence.

## Learning points

The most common form of fungal rhinosinusitis is sinonasal fungus ball which usually presents in adults with normal immunity where maxillary sinus being the most involved site [1, 2]. Moreover, there is female predominance and those of older age as demonstrated in previous studies [3]. It is rare in children. Unilateral signs such as nasal obstruction, purulent rhinorrhea and facial pain are the main complaints of patients.

A CT scan of the sinuses is the imaging of choice, showing a hyperdense calcified formation within a hypodense sinus filling with osteosclerosis of the sinus walls. In our case, the conglomeration of the different calcific formations gave this pseudo tumoral appearance (Figure 2). Treatment was surgical, with removal of the fungal bullets. Functional endoscopic sinus surgery is the definitive approach for fungal ball with high success and low morbidity rates [4, 5, 6].



**Figure 1:** Macrocalcifications in the left maxillary and ethmoidal sinus.



**Figure 2:** Dense conglomeration of fungal hyphae.

### Competing interests

None.

### Patient consent

Obtained.

### References

1. X Dufour, C Kauffmann-Lacroix, JC Ferrie, JM Goujon and MH Rodier, et al. (2006) Paranasal sinus fungus ball: epidemiology, clinical features and diagnosis. A retrospective analysis of 173 cases from a single medical center in France 1989-2002. *Med Mycol* 44(1) :61-67.
2. RS Jiang, WC Huang, KL Liang (2018) Characteristics of sinus fungus ball: a unique form of rhinosinusitis. *Clinical. Medicine Insights: Ear Nose and Throat*.
3. YH Yoon, J Xu, SK Park, JH Heo and YM Kim, et al. (2017) A retrospective analysis of 538 sinonasal fungus ball cases treated at a single tertiary medical center in Korea (1996-2015). *International forum of allergy & rhinology* 7(11) :1070-1075.
4. P Grosjean, R Weber (2007) Fungus balls of the paranasal sinuses: a review *European Archives of Oto-Rhino-Laryngology*. 264 (5) : 461-470.
5. RD DeShazo, K Chapin, RE Swain (1997) Fungal sinusitis. *New England Journal of Medicine* 337( 4): 254-259.
6. MY Seo, SH Lee, G Ryu (2019) Clinical pattern of fungal balls in the paranasal sinuses: our experience with 70 patients. *European Archives of Oto-Rhino-Laryngology* 276 (4): 1035-1038.




This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

[Submit Article](#)

DOI: [10.32474/SJO.2024.10.000339](https://doi.org/10.32474/SJO.2024.10.000339)



### Scholarly Journal of Otolaryngology

#### Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles