



# Self-Medication Patterns Among Medical Students in North India

Neelam Kotwal<sup>1</sup>, Sunil Kumar<sup>2</sup>, Monika Malhotra<sup>2</sup>, Saurabh Sadotra<sup>2</sup>, Pankaj Kumar<sup>3</sup> and Mohammad Sarwar Mir<sup>3\*</sup>

<sup>1</sup>Former Medical Officer at Shri Mata Vaishno Devi Shrine Board, India

<sup>2</sup>IMO Grade 1 ESIC Hospital, India

<sup>3</sup>SKIMS Soura, India

\*Corresponding author: Mohammad Sarwar Mir, SKIMS Soura, India

Received: 📅 December 12, 2018

Published: 📅 December 18, 2018

## Abstract

**Introduction:** Self-medication results in wastage of resources, increases resistance of pathogens and generally causes serious health hazards such as adverse drug reactions, prolonged suffering and drug dependence.

**Method:** A cross-sectional descriptive study was conducted. The participants were medical students from first to final year. The data was collected using a questionnaire. The data was analyzed using SPSS version 2.0.

**Results:** A total of 100 students, 61 (61.00%) male and 39 (39.00%) female, were included in the study. Of the medical students surveyed, self-medication was reported among 88%. The most common ailments for which self-medication were used were: the common cold and headache. The students consulted their textbooks and peers for the medications. Antipyretics and analgesics were the most common self-medicated drugs.

**Conclusion:** The prevalence of self-medication among medical students is high, facilitated by the easy availability of drugs and information from textbooks or seniors. The potential problems of self-medication should be emphasized to the students.

**Keywords:** Self-medication; Medical students; Drugs

## Introduction

Self-medication can be defined as obtaining and consuming drugs without the advice of a physician for diagnosis, prescription or surveillance of treatment [1-3]. Self-medication differs from self-care in that it involves drugs that may do good or cause harm [4]. It has been found that inappropriate self-medication causes wastage of resources, increases resistance of pathogens and generally causes serious health hazards such as adverse drug reactions, prolonged suffering and drug dependence [4-7].

## Material and Methods

A cross-sectional study was undertaken. The study population consisted of medical students from first to final year, within the age group of 18-25 years. The information pertaining to the pattern of self-medication, indications for self-medication and drugs used for

self-medication were included in the questionnaire. The data was analyzed using SPSS version 20.00.

## Results

A total of 100 students participated in the study, of whom 61(61.00%) were male and 31 (39.00%) were female. A total of 88 (88.00%) participants practiced self-medication (Figure 1).

## Discussion

Self-medication is becoming an increasingly important area within healthcare, and this study has shown that it is even more prevalent among medical students. This study has found a prevalence of self-medication of 88 % in medical students in Figure 2 contrast to 59% in a non-medical population in a previous study

[2]. It is also noted that a high level of education and professional status are predictive factors for self-medication [4]. This is similar to the findings in a study conducted by Relend Hem and colleagues (90%) but is higher than the findings (60%) in the study conducted by Henry James and colleagues [4]. In the study it was noticed that the classes of drugs that were commonly used were antipyretics, analgesics, antihistamines and antibiotics. This is similar to studies

done earlier [4,5]. The study has found that self-medication is very common among medical students, facilitated by the easy availability of drugs, and information from textbooks/seniors (Figure 3). Since inappropriate self-medication has the potential to cause serious harm, not only to the students themselves but also to those whom they suggest medication, potential problems of self-medication should be emphasized to the students to minimize this risk.




Figure 1: Conditions prompting self-medication.




Figure 2: Class of drugs for self-medication.




Figure 3: Common sources of information for self-medication.

## References


1. Badiger S, Kundapur R, Jain A, Kumar A, Patanashetty S, et al. (2012) Self-medication patterns among medical students in South India. *AMJ* 5(4): 217-220.
2. Shankar PR, Partha P, Shenoy N (2002) Self-medication and non- doctor prescription practices in Pokhara valley, Western Nepal: a questionnaire based study. *BMC Fam Pract* 3: 17.
3. Montastruc JL, Bagheri H, Geraud T, Lapeyre Mestre M (1997) Pharmacovigilance of self-medication. *Therapie* 52(2): 105-110.
4. James H, Handu SS, Khalid AJ, Khaja A, Otoom S, et al. (2006) Evaluation of the knowledge, attitude and practice of self-medication among first-year medical students. *Med Princ Pract* 15(4): 270-275.
5. Hughes CM, McElnay JC, Fleming GF (2001) Benefits and risks of self-medication. *Drug Saf* 24(14): 1027-1037.
6. Hem E, Stokke G, Reidar Tyssen R, Grønvold NT, Vaglum P, et al. (2005) Self-prescribing among young Norwegian doctors: a nine-year follow-up study of a nationwide sample. *BMC Med* 3: 16.
7. Kiyangi KS, Lauwo JAK (1993) Drugs in home: danger and waste. *World Health Forum* 14(4): 381-384.
8. Sarahroodi S, Arzi A, Sawalha AF, Ashtarinezhad A (2010) Antibiotic Self-Medication among South Iranian University Students. *International Journal of Pharmacology* 6(1): 48-52



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)

DOI: [10.32474/SJPBS.2018.01.000125](https://doi.org/10.32474/SJPBS.2018.01.000125)



### Scholarly Journal of Psychology and Behavioral Sciences

#### Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles