

Can Cats Cause Schizophrenia? an Insight into the Role of Bartonella Henselae in Neuropsychiatric Disorders

Divya R*, Ashok V and Rajajeyakumar M

Department of Physiology, Faculty of medical sciences and research, India

*Corresponding author: Divya R, Department of Physiology, Karpagam Faculty of medical sciences and research, Coimbatore, India

Received: 📅 April 08, 2019

Published: 📅 April 26, 2019

Cat Scratch Disease(CSD)

Bartonella henselae is a fastidious gram-negative rod which is the etiological agent of Cat Scratch Disease(CSD) and is allied with bacillary angiomatosis in HIV-infected individuals. A total of 30 different species of *Bartonella* and 13 of those have been found to infect human beings [1]. Different *Bartonella* species causes wide-ranging and multifaceted diseases in humans [2]. The primary carriers of *Bartonella henselae* are Cats. Stray cats as well as healthy pets. The flea vector *Ctenocephalides felis* transmits *Bartonella* between cats. The mode of transmission to humans is through a scratch or bite from a cat or a kitten. Cat scratch disease is prevalent in all age groups, but more common in children less than 10 years of age [3].

Neurobartonellosis

In Humans, the bacterium infects pericytes, macrophages, erythrocytes, and endothelial cells, resulting in vascular pathology. Throughout the world, research data demonstrates a broad spectrum of neurological abnormalities caused by *Bartonella* spp referred to as neurobartonellosis. It is found to have equal prevalence in immunocompetent and immunocompromised patients [4]. *Bartonella* species were isolated from chronically ill patients, suffering from various diseases such as epilepsy, rheumatoid arthritis and migrane with idiopathic etiology [1].

Pediatric Acute-Onset Neuropsychiatric Syndrome (PANS): A case Study

Due to a sudden onset psychotic behaviour, an adolescent patient was diagnosed and treated for schizophrenia over a period

of 18 months; resulting in failure of psychosis and autoimmune disorders treatment. The lesions on the patient's skin were identified and led to the diagnosis of serum positive *Bartonella* infection. The adolescent was then treated with Long term antimicrobial chemotherapy that helped in full recovery from all the Neuropsychiatric Symptoms. This study advocates the contribution of *Bartonella* infection towards the development and progression of neuropsychiatric disorders such as schizophrenia [1].

Conclusion

This advancement has led to a different aspect of approach towards the infectious etiology of neuropsychiatric illnesses such as Schizophrenia, Alzheimer's disease. Future investigations are essential to understand the potential role of viral and bacterial infections in medically complex neuropsychiatric disorders with idiopathic etiology.

References

1. Breitschwerdt EB, Greenberg R, Maggi RG, Mozayani BR, Lewis A, et al. (2019) *Bartonella henselae* Bloodstream Infection in a Boy With Pediatric Acute-Onset Neuropsychiatric Syndrome. *J Cent Nerv Syst Dis* 11.
2. Vayssier Taussat M, Moutailler S, Féménia F (2016) Identification of Novel Zoonotic Activity of *Bartonella* spp., France. *Emerg Infect Dis* 22(3): 457-462.
3. Mogollon P, Elin (2009) *Bartonella*: emerging pathogen or emerging awareness? *International Journal of Infectious Diseases* 13(1): 3-8.
4. Balakrishnan N, Ericson M, Maggi R, Breitschwerdt EB (2016) Vasculitis, cerebral infarction and persistent *Bartonella henselae* infection in a child. *Parasit Vectors* 9(1): 254.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)

DOI: [10.32474/RRHOAJ.2019.03.000162](https://doi.org/10.32474/RRHOAJ.2019.03.000162)



Research and Reviews on Healthcare: Open Access Journal

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles