



Why to Not Rely on Nature?

Shah Murad^{1*}, Akbar², Salman Iftikhar³, Abdul Ghaffar⁴, Jamila Shah Murad⁵ and Ghazi Mahmood⁶

¹Department of Pharmacology, Islamabad Medical College, Islamabad, Pakistan

²Niazi Teaching Hospital and IMDC, Islamabad Seema, Gynecologist at NMC, Karachi Pakistan

³Department Pharmacology at RLMDC Lahore Pakistan

⁴SWO at LMDC/DANTH Islamabad Pakistan

⁵CI Psychologist at Bahria University, Pakistan

⁶ENT Surgeon at MMC Multan

*Corresponding author: Shah Murad, Department of Pharmacology IMDC, Islamabad-Pakistan

Received:  July 23, 2019

Published:  September 09, 2019

Abstract

Metabolism is the process your body uses to make energy from the food you eat. Food is made up of proteins, carbohydrates, and fats. Chemicals in your digestive system (enzymes) break the food parts down into sugars and acids, your body's fuel. Your body can use this fuel right away, or it can store the energy in your body tissues. If you have a metabolic disorder, something goes wrong with this process. Dyslipidemia is main etiological factor leading to develop coronary artery disease (CAD). Allopathic drugs used in cure of hyperlipidemia, have unwanted effects, so have poor compliance. Now a day's nutraceuticals are getting popularity due to their moderate hypolipidemic actions with fewer adverse effects. Niacin or vitamin B-3 is used as hypolipidemic agent which increases HDL-cholesterol and decreases LDL-cholesterol, VLDL, TGs by different mechanisms. But its main adverse effects are flushing due to synthesis of Prostaglandin D-2 which has vasodilatory effect causing flushing. Cardamom is herb having hypolipidemic potential if used in specific concentration for long time. In this work we did try to compare these two medicines hypolipidemic effects. Study was single blind placebo-controlled conducted at Jinnah Hospital Lahore-Pakistan from July to November 2018. Seventy-five hyperlipidemic patients were selected and divided in three groups. Their base line lipid profile was determined at laboratory of the hospital. Patients were divided in three groups. Group-1 was on placebo therapy, Group-II was on Niacin 1.5 grams daily in divided doses, and Group-III was on Cardamom 1 gram daily in three divided doses. It was two months therapeutic design. At completion of therapeutic regimen after two months we measured all patient's lipid profile. When results were compiled and pre and post treatment values were analyzed statistically, it was observed that both drugs hypolipidemic potential is different, although both have hypolipidemic characteristic, but Niacin is the best for treating Hyperlipidemia. For statistical analysis we used SPSS version 5.0 and paired 't' test was applied to understand significant changes in two tested group's lipid profile.

Introduction

Lipid metabolism disorders, such as Gaucher disease and Tay-Sachs disease, involve lipids. Lipids are fats or fat-like substances. They include oils, fatty acids, waxes, and cholesterol. If you have one of these disorders, you may not have enough enzymes to break down lipids. Or the enzymes may not work properly, and your body can't convert the fats into energy. They cause a harmful amount of lipids to build up in your body. Over time, that can damage your cells and tissues, especially in the brain, peripheral nervous system, liver, spleen, and bone marrow. Many of these disorders can be very serious, or sometimes even fatal. High LDL-cholesterol and low HDL-cholesterol in blood are cause of synthesis of atherosclerotic plaques which get deposited with inner endothelium of coronary

arteries, leading to develop coronary artery disease (CAD) [1,2]. Hyperlipidemia either primary or secondary is challenge for scientists of 21st century due to this disease's complications like coronary artery disease, Hypertension, Cardiac Arrhythmias, and Myocardial Infarction [3,4]. Hyperlipidemia causes LDL particle's oxidation, which initiate formation of atherosclerotic plaques leading to development of coronary artery disease. From this single complication of Hyperlipidemia, all major heart diseases are developed leading to morbidity and mortality due to last lethal major heart disease Ventricular Fibrillation [5-8]. Hypolipidemic drugs commonly used include Statins, nicotinic acid, bile acid binding resins and fibric acids, but all have potential for low patient compliance due to wide range of side effects [9].

Niacin increases apolipoprotein A1 levels due to anti catabolic effects resulting in higher reverse cholesterol transport. It also inhibits HDL hepatic uptake, down-regulating production of the cholesterol ester transfer protein gene. Finally, it stimulates the ABC-A1 transporter in monocytes and macrophages and up-regulates peroxisome proliferator-activated receptor γ results in reverse cholesterol transport. It reduces secondary outcomes associated with atherosclerosis, such as low-density lipoprotein cholesterol, very low-density lipoprotein cholesterol, and triglycerides, but increases high density lipoprotein cholesterol [10]. Despite the importance of other cardiovascular risk factors, high HDL was associated with fewer cardiovascular events independent of LDL reduction. Other effects include anti-thrombotic and vascular inflammation, improving endothelial function [11]. To get good drug-patient compliance many health-related modern researchers have started to put their haeling potential for developing alternatives drugs used in primary or secondary Hyperlipidemia. Cardamom or in urdu ILAICHI is one of the hypolipidemic herb, widely encouraged by cardiologists to be used for prevention of atherogenesis, and coronary artery disease [12]. Cardamum's antioxidant effect is well established in many research studies on medicinal herbs. This nutraceutical contains Flavenoids and Phenolic compounds which scavenge Reactive Oxygen Species (ROS), reducing risk of developing CAD. Oral administration of cardamom extracts significantly reduced total cholesterol, high-density lipoprotein (HDL), low-density lipoprotein, and very low-density lipoprotein and triglycerides in HL patients [13].

Patients & Method

Design: It was single blind placebo-controlled study conducted at GENERAL HOSPITAL Lahore from July to November 2018.

No of Patients: Seventy-five hyperlipidemic patients were selected and enrolled for the study. Written, already explained and approved consent was taken from all patients.

Inclusion Criteria: Inclusion criteria were age limit from 18 to

70 years of both gender male and female primary or secondary hyperlipidemic patients.

Exclusion Criteria: Patients suffering from any vital organ disease or their impaired functions were excluded from the study. Alcoholics, cigarette smokers and patients taking regular medicine for their any physical or mental disease were also excluded.

Grouping: Seventy-five patients were divided in three groups, comprising 25 patients in each group. Group-I was on placebo therapy. They were provided capsules containing grinded rice and mixed wheat. They were advised to take one capsule before meal, thrice daily for two months. Group-II patients were advised to take 1.5 grams of Niacin in three divided doses for the period of two months. Group-III were advised to take one-gram grinded green Cardamom powder mixed in black tea, thrice daily after each meal for the period of two months.

Methods Used: At start of study all patients' blood pressure was recorded and kept in their personal file. Lipid profile of all patients was determined by Frei Dewald Method. Total-cholesterol, LDL-cholesterol and HDL-cholesterol were main parameters we required for further calculation of change in these parameters. All patients were advised to visit clinic fortnightly for their follow up. After two months therapy their lipid profile was measured again by same Frei Dewald Method. Blood Pressure of all participants was again measured and compiled for further statistical analysis.

Statistical Analysis: Data were expressed as the mean \pm SEM and paired "t" test was applied to determine statistical significance as the difference. A probability value of <0.05 was considered as non-significant and $P<0.001$ was considered as highly significant change in the results when pre and post-treatment values were compared

Results

Results are shown in Table 1 with detail and all parameter's values and changes in these parameters are self-explanatory.

Table1: Results of pre and post treatment values, measured in mg/dl (tc, ldl-c, hdl-c) & mm of hg (sbp, dbp).

GROUP	Parameter	At day 0	At day-60	Change	% change	p-value
G-I n=25	SBP	145.13 \pm 1.23	142.09 \pm 1.86	3.0	2.1	>0.05
	DBP	99.77 \pm 1.05	98.67 \pm 2.87	1.1	1.1	>0.05
	TC	228.2 \pm 1.99	226.4 \pm 1.23	1.8	0.8	>0.05
	LDL-C	178.4 \pm 1.67	176.5 \pm 1.09	1.9	1.1	>0.05
	HDL-C	40.7 \pm 1.90	40.9 \pm 2.98	0.2	0.5	>0.05
G-II n=22	SBP	137.11 \pm 2.17	129.66 \pm 2.13	7.5	5.5	<0.01
	DBP	90.96 \pm 1.05	85.94 \pm 2.12	5.0	5.5	<0.01
	TC	235.4 \pm 1.11	204.6 \pm 1.99	30.8	13.1	<0.001
	LDL-C	181.1 \pm 2.87	169.0 \pm 2.22	12.1	6.7	<0.01
	HDL-C	43.5 \pm 1.99	49.1 \pm 1.04	5.6	11.4	<0.01

G-III n=24	SBP	132.54±1.91	130.15±1.05	2.4	1.8	>0.05
	DBP	91.65±1.11	88.78±2.81	2.9	3.2	>0.05
	TC	239.0±2.32	231.8±1.33	7.2	3.0	<0.01
	LDL-C	188.8±2.45	180.0±1.95	8.8	4.7	<0.01
	HDL-C	39.6±1.11	44.5±1.55	4.9	11.0	<0.01

Discussion

Nutraceuticals like vitamins, and nutritional substances like fruits, vegetables, and indian spices are getting popularity in medical research regarding their therapeutic characteristics. Vitamin B-3 or niacin, and Cardamom (illaichi) are well known nutraceuticals which have been proved to have hypolipidemic potential. Metabolic syndrome is cluster of complications in lipid, protein and carbohydrate metabolism in human body. Hyperlipidemia is main fraction and part of this complication. High plasma LDL particles are prone to be oxidized leading to synthesis of atherosclerotic plaques which deposit to endothelial wall of arteries including coronary arteries (CAD). To reduce high plasma fats can prevent individual from being vulnerable for development of CAD. In our results two months therapy with Niacin decreased total and LDL cholesterol 13.1 and 6.7 % respectively. Statistically decrease in total cholesterol is highly significant while change in LDL-cholesterol is significant biostatistically. These results match with results of study conducted by Capuzzi DM et al. [14] who agree with our results. They almost saw same changes in LP (lipid profile) of primary and secondary hyperlipidemic patients.

They mentioned effects of blood pressure which was non-significant when they analyzed pre and post treatment values in systolic and diastolic blood pressure. Our results of change in HDL cholesterol also match with results of Alam K et al. [15] who proved 11% increase in HDL-cholesterol which is highly significant change in pre, and post treatment values of the parameter mentioned. Cantarella L et al. [16] proved significant change in systolic blood pressure of hyperlipidemic patients but non-significant effects on diastolic blood pressure. Goto T et al. [17] explained cause of flushing by using Niacin in hypolipidemic doses. They recommended titration of Niacin dose to improve patient drug compliance. Kawaguchi K et al. [18] did research on Cardamom and found phenolic compounds of this herb as antioxidant. They agree with many researcher's idea or viewpoint that phenolics work as scavenger of Reactive Oxygen Species (ROS). Mittal MK et al. [19] proved that hypolipidemic effects of any herb are not as potent as Statins or Niacin. They have encouraged researchers that active ingredients of medicinal herbs should carefully be extracted and utilized efficaciously when you could preserve all for long time. Our research study proved significant changes in total and LDL cholesterol in 24 hyperlipidemic patients, i.e. 7.2 mg/dl reduction in total cholesterol and 8.8 mg/dl decrease in LDL cholesterol. Changes in both parameters are biostatistically significant.

Almost same results were observed by Bruckert, Eric et al. [20] who again said and advised patient not totally to depend on medicinal herbs, for, it is difficult to treat the disease by monotherapy when they start to develop metabolic syndrome and its serious complications. Babu PV et al. [21] has described that too much active ingredients or chemical compounds present in herbal medicines made it (Cardamom) unpopular because unexpected effects may be observed by using large amount of grinded Cardamom. He warned individuals using large amount of drug may harm metabolic pathways of body, producing unexplained metabolites which can harm body tissues. NA Lokan et al. [22], Temokarr Y et al. [23] explained that there are remarkable number of medicinal herbs and other chemical compounds which scavenge ROS (reactive oxygen species) in human body, preventing development of CAD.

References

1. Reeka VS, Neuendorf HA, Roth JL (2008) Nutrition protocols for the prevention of cardiovascular disease. *Nutr Clin Pract* 23: 468-76.
2. Jagarnath T, Maluava YB (2015) Adipose tissue, inflammation and atherosclerosis. *Journal of atherosclerosis and thrombosis* 20(1): 466-9.
3. Shah M, Ghaffar A, Niaz K, Sheikh AS (2014) ROS and Dyslipidemia. *Ethan Bot J* 12(3): 321-326.
4. Gidda RT, Semonja VC, Faru KK (2014) What about cure of dyslipidemia? *J Cardiovasc Pharmacol* 17: 19.
5. Iyer A, Panchal S, Poudyal H, Brown L (2009) Potential health benefits of Indian spices in the symptoms of the metabolic syndrome: a review. *Indian J Biochem Biophys* 46: 467-481.
6. Kunitomo M (2007) Oxidative stress and atherosclerosis. *Yakugaku Zasshi* 127(12): 199-206.
7. Khitan Z, Kim DH (2013) Fructose: A key factor in the development of metabolic syndrome and hypertension. *J. Nutr. Metab* 13: 1-12.
8. Jia G, Aroor AR, Whaley Connell AT, Sowers JR (2014) Fructose and uric acid: Is there a role in endothelial function? *Curr. Hypertens Rep* 16: 434.
9. Poudyal H, Panchal SK, Waanders J, Ward L, Brown L (2011) Lipid redistribution and metabolism. *J Cardiol* 126(87): 28-30.
10. Tappy L, Le KA, Tran C, Paquot N (2010) Niacin is used as hypolipidemic agent. *Nutrition* 26: 1044-1049.
11. Gass JD (2013) Primary Hyperlipidemia can be treated well by Vitamin B-3 (Niacin). *Lipid Res J* 23(6): 500-510.
12. Verstraeten SV, Fraga CG, Galleano M, Oteiza PI (2010) Basic biochemical mechanisms behind the health benefits of cardamom. *Mol Aspects Med* 31: 435-45.
13. Jenner A, Halliwell B, Rafter J (2010) Use of Cardamom in dyslipidemia. *Am J Clin Nutr* 81: 268-276.


14. Capuzzi DM, Morgan JM, Brusco OA, Intenzo CM (2010) "Niacin dosing: relationship to benefits and adverse effects". *Curr Atheroscler Rep* 2(1): 64-71.
15. Alam K, Pathak D, Ansari SH (2011) Evaluation of Antihyperlipidemic activity of green cardamom. *Int J Pharm Sci Drug Res* 3: 35-37.
16. Cantarella L, Gallifuoco A, Malandra A, Martínková L, Spera A, et al. (2011) Hypolipidemic and Hypotensive effects of Niacin. *Enzyme and Microbial Technology* 48(4): 345-350.
17. Goto T, Teraminami A, Lee JY, Ohyama K, Funakoshi K, et al. (2012) Niacin is good hypolipidemic effects, but it causes flushing. *J Nutr Biochem* 23: 768-776.
18. Kawaguchi K, Mizuno T, Aida K, Uchino K (2011) Green Cardamom and blood lipids. *Biosci Biotechnol Biochem* 61: 102-104.
19. Mittal MK, Florin T, Perrone J, Delgado JH, Osterhoudt KC (2010) 'Use of niacin to beat high plasma lipids as compared to hypolipidemic herbs. *Ann Emerg Med* 50(5): 587-590.
20. Bruckert, Eric, Labreuche, Julien, Amarenco, Pierre (2010) Herbs have hypolipidemic effects, but their wide range of pharmacological effects should be considered. *Atherosclerosis* 21(2): 353-361.
21. Babu PV, Liu D, Hubbarad LR, Kulin MD (2008) Hypolipidemic characteristics of Green Cardamom. *Curr Med Chem* 15: 1840-1850.
22. NA Lokan, HH, Julalu, and TT Tulsik (2015) Do use nutraceuticals for dyslipidemia. *J Thromb* 29(7): 677-679.
23. Temokarr Y, Hulasa V, Fusadaj B (2014) free radical scavenger chemicals found in medicinal plants. *Der J* 6(6): 40-47.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)

DOI: [10.32474/RRHOAJ.2019.04.000176](https://doi.org/10.32474/RRHOAJ.2019.04.000176)



Research and Reviews on Healthcare: Open Access Journal

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles