

Emergence of Pandemics: where we Stand?

Zafar H Tanveer¹, Sajida Zafar², Ali Abuzar Raza³, Rabia Zafar⁴, Jamil Ahmed Lakhair⁵, Shah Murad^{6*}, Shaheena⁷ and Imtenan Sharif⁸

¹HOD Physiology and Principal, QIMS, Quetta Pakistan

²Assistant Professor of Physiology at Nishtar Medical University, Multan Pakistan

³Pathology Department at Nishtar Medical University, Multan Pakistan

⁴DPT Scholar at Rapha International University, Lahore Pakistan

⁵Vice Principal, KIMS, Malir Cantt Karachi Pakistan

⁶Professor of Pharmacology, QIMS, Quetta, Pakistan

⁷Prof of Biochemistry, KIMS, Karachi Pakistan

⁸AP Community Medicine at QIMS Quetta Pakistan

***Corresponding author:** Shah Murad Mastoi, Professor, Department of Pharmacology, QIMS, Quetta, Pakistan

Received:  May 08, 2023

Published:  June 06, 2023

Introduction

Governments around the world, as a matter of policy, are requiring people to socially isolate to mitigate the spread of the novel corona virus. Although an effective tool, this measure carries risk and has brought about huge influence on physically physical and mental health [1]. They are alone-with no certain end to the isolation in view. This study provides evidence for substantial psychological and mental morbidity including complaints deeply human of obsessive-compulsive disorder (OCD) and aggression in the patients who suffered from COVID-19 infection [2]. The new coronavirus outbreak was characterized as a pandemic by the World Health Organization in March 2020. Robust public health containment measures have been widely implemented to prevent transmission of the virus among the global population. The impact of the COVID-19 crisis on population mental health remains unclear, and it is important that the mental health burden of this pandemic on the population is evaluated [3]. Although the frequency of mental disorders is expected to increase during pandemics, the peculiarities of the COVID-19 pandemic can directly impact the clinical course of obsessive compulsive-disorder (OCD), a condition that affects approximately 3% of the general population [4]. OCD is characterized by the presence of obsessions (which are unwanted and unpleasant thoughts, images, or urges) and/or compulsions (repetitive behaviors or mental rituals aimed at reducing the distress provoked by obsessions) [3]. Cleanliness, contamination, and fear of contracting a disease, which are topics of concern for patients with OCD, have now become a

central theme of news and social media. Thus, we would like to elaborate on some possible implications of the COVID-19 pandemic for both diagnosis and clinical decision making about OCD [5].

According to the current biopsychosocial model of psychiatric disorders, socio-cultural, biological, and psychological factors interact synergistically to determine the onset of different disorders. These factors include the patient's beliefs and behaviors, which may be influenced by historical and environmental changes [6,7]. Frequent handwashing, which was previously considered excessive and one of the most common symptoms of OCD, has now been normalized [8]. Currently, considering time-consuming cleaning rituals as a single symptom seems insufficient to diagnose OCD. In such cases, clinical reasoning should counterbalance the degree of protection these rituals produce with their level of interference in functioning. Such reasoning is required to confirm an OCD diagnosis or consider treatment changes for those already on medication or in cognitive-behavioral therapy [9]. Patients may experience a worsening of OCD symptoms in different dimensions, including, but not limited to, contamination/cleaning, aggression, and hoarding dimensions. Patients who had never presented such symptoms may experience their onset in the context of this major environmental change. Higher levels of avoidant behavior are also expected. Moreover, OCD patients can experience a worsening of depression and anxiety symptoms during major life events, among which the current pandemic should be included [10].

Exposure and ritual prevention, a key behavioral technique in OCD treatment, should be carefully tailored during this period. Recommending unrestricted exposure to feared stimuli may prove imprudent. Psychological strategies for treating OCD should consider the well-being and safety of patients. Pharmacological strategies should be guided by the best evidence-based recommendations [11].

Protection recommendations (e.g., hand washing) can reinforce the irrational beliefs of patients with OCD and poor insight. Therefore, engagement in exposure and ritual prevention activities may be lower, which could impact the long-term prognosis for OCD [12]. We assumed that patients and their relatives would show similar increased levels of anxiety, and that anxiety levels would be higher than depressive levels. This hypothesis is based upon the unpredictable nature of the COVID-19 and the accompanying uncertainty regarding the course of the illness and its infectious potential, which are key factors for anxiety [13]. Among relatives, we hypothesized that children would show decreased anxiety and depression levels, compared to adult relatives. This hypothesis is based on the notion that although children must deal with the same negative feeling of anxiety as adults, they do not share the same objective burdens as adults, such as caring for family functions at this difficult time [14]. This paper aims specifically to investigate the mental status of those patients who have recovered from COVID-19 pandemic. Understanding the impact of self-isolation & self-quarantine on mental health will help us mitigate these mental disturbances through appropriate developments in mental health services such as online consultation and digital psychiatric management to keep the patient's mental health in check.

Methodology

Study Design: Our Study is cross sectional study.

Study Settings: Study is carried out among those patients which were isolated for COVID19.

Study Duration: The study has been conducted for 6 months from March 2021 to August 2021.

Study Population: A total of 320 patients completed this study through the online Google forms.

Sampling technique: Non - Probability Convenient Sampling

Sample Size: Sample size $n = 320$ Where $n =$ sample size $z = z$ value corresponding to a given confidence level = 1.96 $p =$ % frequency of the outcome factor = 58 % $c =$ standard error/ effect size $n = 320$.

Inclusion criteria: 1) those patients which are positive for COVID 19. 2) isolate for COVID 19. 3) Should go through the symptoms of anxiety and OCD.

Exclusion criteria: 1) those patients which were admitted to the hospital. 2) Any previous history of psychiatry disorders. 3) those which are not isolated for COVID19.

Data collection: Data is collected through a carefully structured Google form for post isolated COVID19 patients. Patients who volunteered were asked to fill in the questionnaire and demographic Performa after informed consent was obtained from each one of them. The participants could choose multiple reasons applicable to themselves. Data was collected regarding symptoms and severity of anxiety and obsession compulsive disorders (OCD) in COVID 19 patients after isolation. For data collection procedure regarding anxiety, we used Hamilton Anxiety Rating Scale (HAM-A) and for obsession compulsive disorders (OCD) we used Yale - Brown scale.

Hamilton Anxiety Rating Scale (HAM-A). The HAM-A was one of the first rating scales developed to measure the severity of anxiety symptoms and is still widely used today in both clinical and research settings. The scale consists of 14 items, each defined by a series of symptoms, and measures both psychic anxiety (mental agitation and psychological distress) and somatic anxiety. The scale consists of 14 items, each defined by a series of symptoms, and measures both psychic anxiety (mental agitation and psychological distress) and somatic anxiety (physical complaints related to anxiety). Although the HAM-A remains widely used as an outcome measure in clinical trials, it has been criticized for its sometimes-poor ability to discriminate between anxiolytic and antidepressant effects, and somatic anxiety versus somatic side effects. Each item is scored on a scale of 0(not present) to 4 (severe), with a total score range of 0-56, where greater than 17 indicates mild severity, 18-24 mild to moderate severity and 25-30 moderate to severe. The Yale-Brown Obsessive-Compulsive Scale is a standardized rating scale with both clinician administered and self-report versions available, measuring 10 items pertaining to obsessions and compulsions on a five-point Likert scale. Scores range from 0 (no symptoms) to 4 (severe symptoms), and a total score is calculated by summing items 1 to 10 and can range from 0 to 40.

Ethical consideration: Was also taken in account throughout the process of the study. Data collection process was started after approval from the Ethical Review Committee of Karachi Institute of Medical sciences (KIMS).

Data Analysis: The data were entered and analyzed on SPSS V26.0. Frequency and percentages tables were presented.

Results

Demographic Characteristics

The study collected data from 320 patients. Analysis of data collected about patients is. $N=320$ patients involved in the survey, Females $N=113$ (35.4%) whereas $N=207$ (64.6%) were males. Patient's qualifications, results demonstrated that (3.7%) patients had a primary education, (1.8%) patients had a middle education, (3.7%) patients had a matric level education, (8.6%) had an intermediate level education, (55.2%) had undergraduate education. qualification, (25.5%) patients had a post graduate qualification. Further age, (75.2%) had age. between 20-30 years,

(10.1%) had age between 30-40 years, (5.2%) had age between 40-50 years, (7.7%) had aged between 50-60 years. Data on current employment status revealed that there were (62.3%) patients with current employment and (35.9%) patients were without employment. Patients' marital status, (70.9%) patients were single, (25.8%) patients were married, (6%) patients were divorced. Regarding monthly household income, (40.5%) had below 50,000, 79 (24.2%) had above 50,000, 109 (33.4%) had above 1,00,000. Family structure, (52.5%) had nuclear, (39.9%) had joint, (5.8%) had extended family structure.

Data on methods used for screening Covid-19, (89.0%) were through PCR and (9.2%) were through serology. Data on isolation, (96.0%) were isolated at home and (4%) were isolated at hospital. Further analysis shown in Table 2.1 of the survey data obtained in this project specifically for OCD. and anxiety in post COVID patients shows that gender has no effect on anxiety levels with almost equal ratios for males and females i.e., there are (64.6%) were males and females accounted for (35.4%).

Discussion

Findings from the previous indicate that COVID-19 is eliciting distress in those with OCD, as well as in non-clinical individuals. The two groups considered high and low on the severity of OCD symptoms were significantly different on the severity of COVID-19-related intrusive thoughts, the high group having a greater severity. COVID-19-related intrusions were present in the low OC severity group and were considered mildly distressing. Distress related to intrusive thoughts about COVID-19 was significantly higher than typical OC-related intrusive thoughts for those with a higher OCD severity. Frequency, distress, and difficulty dismissing COVID-19-related intrusive thoughts were all significantly associated with one's OCI-R severity (excluding distress for the low severity group), indicating that the severity of one's OCD symptoms was related to the severity of COVID-19 intrusive thoughts. Also, the severity of COVID-19-related intrusive thoughts was maintained over time in both the high and low OC severity groups. The previous findings indicate that individuals with a higher OC severity were more impacted by the presence of COVID-19 intrusive thoughts. Specifically, their COVID-19 intrusive thoughts were significantly more distressing than their typical OC intrusive thoughts. For the entire sample, the higher severity of OC intrusive thoughts, the higher the severity of COVID-19 related intrusive thoughts.

From a cognitive-behavioral perspective, maladaptive beliefs regarding risk and one's ability to cope influence interpretations of physical symptoms, intrusive thoughts, or social interactions, thereby contributing to elevated anxiety. Anxiety occurs when we perceive threats, and individuals with anxiety disorder diagnoses tend to perceive threats as more likely or costly than is objectively the case. However, the same processes would be involved in the case of a real threat. Specifically, substantial distress can also occur from focusing on the aspects of a real threat that

one perceives as unchangeable. Indeed, finding a healthy balance between addressing a threat versus suppressing thoughts about it can be difficult. In some cases, the decreases in anxiety from avoidance can tip the balance too far, strengthening beliefs regarding danger and the need to avoid. While previous studies have recently been published on OCD in clinical populations during the pandemic [1], the presence of intrusive thoughts, or obsessions, in the general population has not been studied to our knowledge at this point. The previous study only included students enrolled in undergraduate or graduate studies at a state university with a limited sample size. However, young adults have been identified as an at-risk population for mental health issues specifically during this pandemic [2]. The previous study investigates psychopathology in a sample of COVID-19 survivors at one month follow-up after hospital treatment [3]. They reported high rates of PTSD, depression, anxiety, insomnia, and OC symptomatology. The findings of the psychiatric morbidities ranged from 10% to 35% in the post-illness stage [4] With regard to the risk factor related to psychopathology, consistently with previous epidemiological studies have found that females, and patients with positive previous psychiatric diagnoses, suffered more in all psychopathological dimensions [5]. Moreover, outpatients showed increased anxiety and sleep disturbances, while the duration of hospitalization inversely correlated with PTSD, depression, anxiety, and OC symptomatology. Also considering the worse severity of COVID-19 in hospitalized patients, this observation suggests that less healthcare support could have increased the social isolation and loneliness typical of COVID-19 pandemics, thus inducing more psychopathology after remission [6]. Finally, younger patients showed higher levels of depression and sleep disturbances, in agreement with previous studies describing a worse psychological impact of COVID-19 pandemic in younger people [7].

Neither oxygen saturation level at follows up nor baseline inflammatory markers associated with depression, anxiety, PTSD nor insomnia, suggesting that psychiatric symptomatology was not a manifestation of physical symptoms, except for baseline SII that positively associated with measures of anxiety and depression at follow-up. The SII is an objective marker of the balance between host systemic inflammation and immune response status considering together neutrophil, platelet, and lymphocyte all of them involved in different pathway of immune/inflammatory response [8]. Higher levels have been associated with worse prognosis in several medical diseases, in the field of oncology. In a single study, higher SII levels were associated with major depressive disorder (MDD) [9], suggesting that it could be a marker of the low-grade inflammation observed in mood disorders [10]. Recent evidence, in agreement with our observation, suggests an impact of COVID-19 on OCD related. In the light of the above, interest to deepen research on biomarkers of inflammation is warranted, to investigate the possible association between possible persistent low-grade inflammation as observed in mood disorders [11], and psychopathological symptoms at follow-up in COVID-19 survivors. This approach could also allow us to identify

possible new specific targets for the treatment of inflammation-related neuropsychiatric conditions. According to our study conducted in post-isolated COVID 19 patients, it is reported that there are high rates of PTSD, depression, anxiety, insomnia, and OC symptomatology.

Our findings from Hamilton scale A, Alpha value is 0.919 % and from the previous studies Alpha value was = 0.75 %. Where mean Male (25.23), and female (28.34) which shows there is difference in level of anxiety in both genders. Psychiatric consequences to SARS-CoV-2 infection can be caused both, by the immune response to the virus itself, or by psychological stressors such as social isolation, psychological impact of a novel severe and potentially fatal illness, concerns about infecting others, and stigma. The immune response to coronaviruses induces local and systemic production of cytokines, chemokines, and other inflammatory mediators [12]. COVID-19 patients, such as SARS and MERS patients, show high levels of Interleukin (IL)-1 β , IL-6, Interferon (IFN)- γ , CXCL10, and CCL2 suggesting an activation of T-helper-1 cell function. Moreover, in COVID-19, unlike in SARS and MERS, elevated levels of T-helper-2 cell-secreted cytokines (such as IL-4 and IL-10) were found [13]. Higher concentrations of these cytokines seem to suggest a more severe clinical course [14]. Cytokines dysregulation (especially IL-1 β , IL-6, IL-10, IFN- γ , TNF- α , and transforming growth factor- β (TGF- β)) are known to involve factors that others and we associated with psychiatric disorders [15]. Neuroinflammation, blood-brain-barrier disruption, peripheral immune cell invasion into the CNS, neurotransmission impairment, hypothalamic-pituitary-adrenal (HPA) axis dysfunction, microglia activation and indoleamine 2,3-dioxygenase (IDO) induction, all represent interaction pathways between immune systems and psychopathological mechanism underpinning psychiatric disorders [16]. However, the findings of the current study and related work are encouraging that we may have tools to alleviate some of the COVID-related anxiety and distress [16-42].

References

- Pfefferbaum B, North CS (2020) Mental Health and the Covid-19 Pandemic. *N Engl J Med* 383(6): 510-512.
- Silva RM, Shavitt RG, Costa DL (2020) obsessive-compulsive disorder during the COVID-19 pandemic. *Braz J Psychiatry* 43(1): 108.
- Shavitt RG, de Mathis MA, Oki F, Ferrao YA, Fontenelle LF, Torres AR, et al. (2014) Phenomenology of OCD: Lessons from a large multicenter study and implications for ICD-11. *Journal of Psychiatric Research* 57: 141-148.
- Engel G (1980) The clinical application of the biopsychosocial model. *Am J Psychiatry* 137(5): 535-544.
- Stein DJ, Costa DLC, Lochner C, Miguel EC, Reddy YCJ, et al. (2019) Obsessive- compulsive disorder. *Nat rev Dis primers* 5(1): 52.
- Grillon C, Lissek S, Rabin S, McDowell D, Dvir S, et al. (2008) Increased Anxiety During Anticipation of Unpredictable But Not Predictable Aversive Stimuli as a Psychophysiological Marker of Panic Disorder. *Am J Psychiatry* 165(7): 898-904.
- Montgomery RJV, Gonyea JG, Hooyman NR (1985) Caregiving and the Experience of Subjective and Objective Burden. *Family Relations* 34(1): 19.
- Abbing A, Baars EW, de Sonnevile, Leo, Ponstein AS, et al. (2019) The effectiveness of art therapy for anxiety in adult women: A randomized controlled trial. *Front Psychol* 10: 1203.
- Abrams MP, Wan EY, Waase MP, Morrow JP, Dizon JM, et al. (2020) Clinical and cardiac characteristics of COVID-19 mortalities in a diverse New York City Cohort. *J Cardiovasc Electrophysiol* 31(12): 3086-3096.
- Acuña Castroviejo D, Escames G, Figueira JC, de la Oliva P, Borobia AM, et al. (2020) Clinical trial to test the efficacy of melatonin in COVID-19. 69(3): e12683.
- Adell A (2010) Lu-AA21004, a multimodal serotonergic agent, for the potential treatment of depression and anxiety. *IDrugs* 13(12):900-910.
- Ahmed H, Patel K, Greenwood D, Halpin S, Lewthwaite P, et al. (2020) Long-term clinical outcomes in survivors of severe acute respiratory syndrome and Middle East respiratory syndrome coronavirus outbreaks after hospitalisation or ICU admission: A systematic review and meta-analysis. *J Rehabil Med* 52(5): jrm00063.
- Al Sarraj S, Troakes C, Hanley B, Osborn M, Richardson MP, et al. (2020) Invited Review: The spectrum of neuropathology in COVID-19. *Neuropathology and Applied Neurobiology* 47(3): 3-6.
- Lima KA de, Rustenhoven J, Mesquita SD, Wall M, Salvador AF et al. (2020) Meningeal $\gamma\delta$ T cells regulate anxiety-like behavior via IL-17a signaling in neurons. *Nat Immunol* 21(11): 1421-1429.
- Vahia V (2013) Diagnostic and statistical manual of mental disorders 5: A quick glance. *Indian J Psychiatry* 55(3): 220-223.
- Awasthi S, Pan H, LeDoux JE, Cloitre M, Altemus M, et al. (2020) The bed nucleus of the stria terminalis and functionally linked neurocircuitry modulate emotion processing and HPA axis dysfunction in posttraumatic stress disorder. *NeuroImageClin* 28:102442.
- Baker A, Simon N, Keshaviah A, Farabaugh A, Deckersbach T, et al. (2019) Anxiety Symptoms Questionnaire (ASQ): development and validation. *Gen Psychiatr* 32(6).
- Baptista AF, Baltar A, Okano AH, Moreira A, Campos ACP, et al. (2020) Applications of Non-invasive Neuromodulation for the Management of Disorders Related to COVID-19. *Front Neurol* 11: 573718.
- Bate J, Malberg N (2020) Containing the Anxieties of Children, Parents and Families from a Distance During the Coronavirus Pandemic. *J Contemp Psychother* 50(4): 284-294.
- Beerse ME, Van Lith T, Stanwood G. (2020) Therapeutic psychological and biological responses to mindfulness-based art therapy. *Stress Health* 36(4): 419-432.
- Behan C (2020) The benefits of meditation and mindfulness practices during times of crisis such as covid-19. *Irish Journal of Psychological Medicine* Pp 1-3.
- Bienvenu OJ, Friedman LA, Colantuoni E, Dinglas VD, Sepulveda KA, et al. (2017) Psychiatric symptoms after acute respiratory distress syndrome: a 5-year longitudinal study. *Intensive Care Med* 44(1): 38-47.
- Bjelland I, Dahl AA, Haug TT, Neckelmann D. (2002) The validity of the Hospital Anxiety and Depression Scale. *Journal of Psychosomatic Research* 52(2):69-77.
- Boehmer TK, DeVies J, Caruso E, Santen, Tang S, Black CL, et al. (2020) Changing Age Distribution of the COVID19 Pandemic United States, May August. *MMWR Morbidity and mortality weekly report* 69(39): 1404-1409.


25. Brodin P (2021) Immune determinants of COVID-19 disease presentation and severity. *Nature Medicine*. 27(1): 28-33.
26. Buckner JD, Bernert RA, Cromer KR, Joiner TE, Schmidt NB. (2008) Social anxiety and insomnia: the mediating role of depressive symptoms. *Depression and Anxiety*. 25(2): 124-130.
27. Buemann B, Marazziti D, Uvnäs-Moberg K (2020) Can intravenous oxytocin infusion counteract hyperinflammation in COVID-19 infected patients? *The World Journal of Biological Psychiatry*. 22(5): 387-398.
28. Casillo GM, Mansour AA, Raucci F, Saviano A, Mascolo N, et al. (2020) Could IL-17 represent a new therapeutic target for the treatment and/or management of COVID-19-related respiratory syndrome? *Pharmacological Research* 156:104791.
29. Alagoz O, Sethi AK, Patterson BW, Churpek M, Alhaneaee G, et al. (2021) The impact of vaccination to control COVID-19 burden in the United States: A simulation modeling approach. *PloS One* 16(7): e0254456.
30. (2020) Coronavirus Disease 2019 (COVID-19). Centers for Disease Control and Prevention.
31. Chakraborty A, Karmakar S (2020) Impact of COVID-19 on obsessive compulsive disorder (OCD). *Iranian Journal of Psychiatry* 15(3): 256-259.
32. Glowacz F, Schmits E. (2020) Uncertainty and Psychological Distress during lockdown during the COVID-19 Pandemic: the young adults most at risk. *Psychiatry Research*. 293: 113486.
33. Lee SA (2020) Coronavirus anxiety scale: A brief mental health screener for COVID-19 related anxiety. *Death Studies* 44(7): 393-401.
34. Mazza MG, De Lorenzo R, Conte C, Poletti S, Vai B, et al. (2020) Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors. *Brain Behavior and Immunity* 89: 594-600.
35. Cameron AC, Gelbach JB, Miller DL (2008) Bootstrap-Based Improvements for Inference with Clustered Errors. *The Review of Economics and Statistics* 90(3): 414-27.
36. Huang C, Wang Y, Li X, Ren L, Zhao J, et al. (2020) Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395(10223): 497-506.
37. Blasco BV, García-Jiménez J, Bodoano I, Gutiérrez-Rojas L (2020) Obesity and Depression: its Prevalence and Influence as a Prognostic Factor: A Systematic Review. *Psychiatry Investigation* 17(8): 715-724.
38. Ozbay F, Johnson DC, Dimoulas E, Morgan CA, Charney D (2007) Social support and resilience to stress: From neurobiology to clinical practice. *Psychiatry (Edgmont)* 4(5): 35-40.
39. Ma Z, Zhao J, Li Y, Chen D, Wang T, et al. (2020) Mental health problems and correlates among 746 217 college students during the coronavirus disease 2019 outbreak in China. *Epidemiology and Psychiatric Sciences* 29: e181.
40. Hou F, Yu T, Du R, Fan G, Liu Y, et al. (2020) Clinical course, and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet* 395(10229): 1054-1062.
41. Mazza MG, De Lorenzo R, Conte C, Poletti S, Vai B, Bollettini I, et al. (2020) Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors. *Brain Behavior and Immunity* 89: 594-600.
42. Benedetti F, Mazza M, Cavalli G, Ciceri F, Dagna L, Rovere-Querini P, et al. (2020) Can Cytokine Blocking Prevent Depression in COVID-19 Survivors? *Journal of Neuroimmune Pharmacology* 16(1): 1-3.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)

DOI: [10.32474/RRHOAJ.2023.08.000279](https://doi.org/10.32474/RRHOAJ.2023.08.000279)



Research and Reviews on Healthcare: Open Access Journal

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles