



Treatment of Lumbar Intradiscal Pathology by Means Of percutaneous Through the Use of Tissue Micrografting Soft of The Auricular Cartilage (Perichondrium), of The Patient Himself

Iñaki Arrotegui*

Department Of Neurosurgery, Spain

*Corresponding author: Iñaki Arrotegui, Department Of Neurosurgery, Spain

Received: 📅 February 15, 2019

Published: 📅 February 26, 2019

Introduction

Microinjerto Tisular: The principle on which this technology is based is to use healthy counterpart connective tissue of the same patient processed with KIT to regenerate its own damaged tissue. The affinity of the donor and recipient tissue used contributes a high differentiation and potentiality obtaining as a result a great cellular regenerative efficiency. Patented technology applied to a surgical instrument, sterile and disposable, which by means of mechanical disintegration in small tissue particles isolates SVFs with high regenerative power. 1: 1 ratio With 1 cm² of healthy tissue (biologically different) We regenerate 1cm² of damaged tissue. Proportion 1:20 With 1 cm² of healthy tissue (biologically homologous) we regenerate 20 cm² of damaged tissue.

New Microfinance Theory: It is based on the Theory of MICROINJERTO TISULAR. Using WOVEN HOMOLOGOUS FABRIC (AUTOLOGOUS) + Kit you get how: THE REGENERATION OF DAMAGED TISSUE. Surgical concept: The smaller the dimension of the graft, the easier it is to integrate into the implanted tissue. Biological concept: The size of the SVF (Vascular Stromal Fracture) cells is approx. 50 microns. This selection and isolation we obtain are those that have a greater potentiality, greater power of differentiation and a high efficiency in cell regeneration. Example: To regenerate articular cartilage we use: Auricular Cartilage + Perichondrium Articular cartilage: It is a cartilage of hyaline type that lacks vascularization whose free surface is not covered by perichondrium. Perichondrium: A layer of fibrous and compact connective tissue that lines the cartilage, except the joint. It is highly innervated and vascularized.

Regenerative Homology

When we extract healthy connective tissue HOMOLOGOUS to the damaged tissue that we want to regenerate, the action of the

SVF induced by the pericytes and endothelial cells that increase its vascularization is increased, therefore we take advantage of 95% of its:

- a) DIFFERENTIATION CAPACITY
- b) POTENTIALITY
- c) EFFICIENCY

When we extract healthy connective tissue DIFFERENT from the damaged tissue that we want to regenerate, only fibrocytes, chondrocytes and osteoblasts would be acting, so WE USE ONLY APROX. 5% of your:

- a) DIFFERENTIATION CAPACITY
- b) POTENTIALITY
- c) EFFICIENCY

The only cells with the capacity to differentiate and therefore with the potential to regenerate are the SVF (Vascular Stricture Fraction). This type of cells represent approximately 5% of the cell population when we perform a Micrograft.

Method and Technique

Cell preparation and intervention in the study

- I. PROTOCOL: PATHOLOGY of COLUMN
 - a. Application of the SVF Injection of ACTIVE REGENERA
 - b. Number of Punch 3 punch of 2,5mm
 - c. How much Serum Injectable 0.2ml per facet
 - d. How many minutes of motor 6 minutes

- e. Extraction area of the Punch Pavilion Auriculares
- f. Pathology of the Column - Facetario Syndrome - Pain.
- g. Where is Injected: nucleus pulposus: Focus of Pain
- h. Special Material -Trocars appropriate for
- i. infiltration of the spine (It must be done in the operating room)

Method

- A. Fill the Rigeneracons kit, if three levels are made with 1.2 ml of injectable physiological saline, per facet is 0.2 ml.
- B. Insert the punch into the kit in the corresponding department under the knife.
- C. Process the indicated time.
- D. Extract the result of the process with a syringe
- E. Application to the area to be treated.

Discussion

Degenerative disc disease is associated with symptoms such as pain and possibly; weakness or numbness of the MMIII. Until recently, patients had few options. Surgery requires extensive recovery, and time out. work, usually at least 6 weeks. In addition, the risk of complications is significant, apart from the complications that may arise from anesthesia, as well as the complications of the intervention, there is a risk of 1 in 10,000 of bowel or bladder incontinence and a risk of 1 in 1000 of nerve root damage. There may also be 1-3% risk of CSF leak, 1% risk of infection and 5-10% risk of spinal instability [1-3]. SVF does not require culture expansion in vitro and it is easy to perform the extraction. These cells can be placed directly on the disc nuclei using a minimally invasive technique guided by fluoroscopy [4,5]. Clinical studies have demonstrated the safety and feasibility of using SVF in patients with degenerative disc. No major safety problems were observed and the procedures were well tolerated in all patients. In addition, patients showed statistically significant improvements in several parameters including flexion, pain classifications, VAS, PPI and questionnaires in abbreviated form. Although ODI and BDI did not show statistically significant changes due to the low number of subjects in the trial, the data allow to verify positive trends. In addition, most patients reported improvements in their Dallas Pain Questionnaire scores [6,7]. Although the study suggests that the

use of SVF is safe and feasible, further studies would be necessary to determine the true clinical effect of the treatment. Given the encouraging results in this small sample size with statistical significance, other clinical studies would be necessary [8]. Several parameters showed statistically significant improvements over a period of 6 months. A true assessment of efficacy and safety would require further phase II / III studies. However, the current study provides encouraging viability data on the intradiscal treatment of stem cells and suggests some clinical benefits of SVF therapy in patients with degenerative disc [9].

Conclusion

The present study will try to define the safety and viability of intradiscal transplantation of autologous SVF in patients with degenerative disc disease and perform the comparison with patients treated by Conservative or Nucleoplasty treatment.

References

1. Miranville A, Heeschen C, Sengenès C, Curat CA, Busse R, et al. (2004) Improvement of postnatal neovascularization by human adipose tissue-derived stem cells. *Circulation* 110(3): 349- 355.
2. Lihua Zhou, Qun Song, Jiangwei Shen, Luwei Xu, Zheng Xu, et al. (2017) Comparison of human adipose stromal vascular fraction and adipose-derived mesenchymal stem cells for the attenuation of acute renal ischemia/reperfusion injury. *www nature com Scientific Reports*.
3. Zain Khalpey (2014) First in Man: Adipose-derived Stromal Vascular Fraction Cells May Promote Restorative Cardiac Function. *The American Journal of Medicine* 127(5):11-12.
4. Ceccarelli G, Gentile P, Marcarelli M, Balli M, Ronzoni F, et al. (2017) In Vitro and In Vivo Studies of Alar-Nasal Cartilage Using Autologous Micro-Grafts: The Use of the Rigenera® Protocol in the Treatment of an Osteochondral Lesion of the Nose. *Pharmaceuticals* 10(2): E53.
5. Rodriguez y Baena R, D Aquino R, Graziano A, Trovato L, Aloise AC, et al. (2017) Auto-logous Periosteum-Derived Micrografts and PLGA/HA Enhance the Bone Formation in Sinus Lift Augmentation. *Front Cell Dev Biol* 5: 87.
6. D Aquino R, Trovato L, Graziano A, Ceccarelli G, De Angelis GC, et al.(2016) Periosteumderived micro-grafts for tissue regeneration of human maxillary bone. *J Transl Sci* 2(2): 125-129
7. Monti M, Graziano A, Rizzo S, Perotti C, Del Fante C, et al. (2016) In Vitro and In Vivo Differentiation of Progenitor Stem Cells Obtained After Mechanical Digestion of Human Dental Pulp. *J Cell Physiol* 232(3): 548-555.
8. Gentile P, Scioli MG, Bielli A, Orlandi A, Cervelli V (2016) A combined use of Chondrocytes Micro Grafts (CMG) Mixed with Platelet Rich Plasma (PRP) in Patients Affected by Pinch Nose Deformity. *J Regen Med* 5(2).
9. Noda S, Sumita Y, Ohba S, Yamamoto H, Asahina I (2017) Soft Tissue Engineering with Micronized-Gingival Connective Tissues. *Journal of Cellular Physiology* 233(1):249-258.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)



Surgery & Case Studies: Open Access Journal

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles