

Survival After Radical Surgery for Huge Sarcomatoid Bladder Tumor

Safarik L^{1*}, Handrejch L², Koch M², Horáková M² and Popílka I³

¹Libor Safarik, Urology Clinic, Czech Republic

²Department of Urology and Surgery, General District Hospital, Czech Republic

³Department of Surgery, NH Hospital Horovice, Czech Republic

*Corresponding author: Safarik L, Urology Clinic, Czech Republic, Czechia

Received:  May 10, 2019

Published:  May 28, 2019

Case Report

A 58-year-old male with BMI 36,8 was seen due to his allegedly three months lasting dysuria and traces of blood in his urine. His renal function was normal, CRP 18,7 and urinary sediment was flooded with red blood cells, but w/o bacterial infection. There was no dilatation on U/S, but hyperechoic exophytic mass was seen in the bladder. The CT scan showed mass, as on Figure 1a. Biopsy from the mass was taken to disclose sarcomatoid tumor of the bladder. (Figure 2a). Since there was no secondaries observable in time of CT scan and on scintigraphy, the radical cystectomy with ileal conduit was suggested. The patient agreed immediately, but there was the cardiac status in the way, since he had hardly survived severe cardiac attack with successive cardiac failure only two years ago, which did not resolved completely. The cardiac output, e.g. ejection fraction (EF), had improved only on permanent pacemaker to reach only 42%. Patient was made aware of possible serious complications after the surgery, cardiac arrest and possibility of re-operation due to his obesity and unfavorable medical status due to general ischemia and atherosclerosis. Despite that, he opted for surgical approach, since the malignancy was undisputable, he was incontinent due to lack of bladder capacity and there was intermittent urinary bleeding. Radical cystoprostatectomy with incontinent Brickere's diversion was carried out showing deep bladder smooth muscle involvement (Figure 3a), but without pelvic nodal infiltration. Moreover, early stage prostate cancer pT1a was found despited unsuspected PSA 1.48ug/L, preoperatively. The patient was discharged on the day 16 and despite trained in stoma care together with his wife, it became very soon clear, that this procedure was out of the capabilities of the couple. The patient did not comply with ambulatory care at all, he was not able to keep the stoma clean and hence the cutaneous plate for collecting urinary bag was difficult to seal on the fatty abdominal wall resulting in urinary leakage around it. Patient had to be re-admitted to the hospital and local therapy for subcuticular abscess around the stoma had to carried out. He was discharged

cured, (Figure 4a) and IVU showed normal collecting systém on both sides and ileal conduit with no urinary leakage, (Figure 5a). One month later, the man was admitted to the hospital again, due to urosepsis. As it was disclosed later on, he attempted to stop the urine coming out of the stoma with plugging it with the swab (!). He lost on weight substantially and surgical repair of the ileal conduit was suggested to cut it short a little bit, since there was no need to have it so long as it was done originally, when the patient suffered from several obesity. He refused it and bilateral nephrostomy was done successively due to urinary leakage from the abscessed conduit. Patient was left on supra-conduit urinary diversion as the permanent urinary diversion and one year after the radical surgery, he was still without detectable metastases. No adjuvant radio- or chemotherapy was done to prevent possible later malignant spread. The patient succumbed to the widespread metastatic disease after another year, accomplishing two full years after diagnosis and 22 months after radical surgery.




Figure 1a.




Figure 2a.




Figure 3a.



Figure 4a.



Figure 5a.

Discussion

Sarcomatoid bladder tumors have very bad reputation regarding the patients' survival and possible treatment. If diagnosed, the afflicted patient has his days usually numbered, as generally observed [1,2]. Despite radical approach and adjuvant chemo and possible radiotherapy, the course of the disease remains straight forward and leads to deaths, almost inevitably. The course of the disease is generally considered to be quick, and the patients experience frequent complications, as recurrent bleeding, dysuria, infection, urosepsis, sometimes also post-renal anuria. In this case, patient was clinically unfit for chemotherapy from the very beginning, but an adjuvant possible radiotherapy on the pelvis minor (to kill the possible micrometastases in the time of surgery) might have proved to be beneficial. However, the major problem after successful surgery was patient's uncompliance with post-operative treatment and stoma care, as well as huge obesity. Despite that, the highly demanding surgery was met with quite unusual long patient's survival, but quality of life was rather dodgy regarding post-operative infections and subsequent urinary drainage through nephrostomic catheters. The patient was quite young, 58, had clinical difficulties with bladder capacity and urinary bleeding. He did not show any mental disorders prior to surgery, but his mental status had probably been underestimated with miserable postoperative compliance with successive medical care and possible oncological adjuvant therapy.

References


1. Lopez-beltran A, Pacelli A, Rothenberg HJ (1998) Carcinosarcoma and sarcomatoid carcinoma of the bladder: clinicopathological study of 41 cases. *J Urol* 159(5): 1497-1503.
2. Uemura K, Kawahara T, Ishida H, Nakaigawa N (2018) Sarcomatoid Variant of Bladder Carcinoma: A Case Report. *Case Rep Oncol* 11(3): 633-637.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)

DOI: [10.32474/JUNS.2019.01.000125](https://doi.org/10.32474/JUNS.2019.01.000125)



Journal of Urology & Nephrology Studies

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles