



Private Health Insurance is a Favourable Prognostic Risk Factor in the Outcomes of Mortality Post Radical Cystectomy

Simon Robinson*, Ailsa Butler, Mufeed Ali and Hanif Motiwala

FHFT Department Urology, Oxon dept epidemiology Oxford, Frimley health foundation trust, UK

*Corresponding author: Simon Robinson, Frimley health foundation trust, FHFT Department Urology, Oxon dept epidemiology Oxford, FHFT Dept Pathology, UK

Received: 📅 July 27, 2020

Published: 📅 August 07, 2020

Abstract

Objectives: There are well recognised differences in health outcomes generally in society due to a range of socio-economic factors. We have compared survival outcomes in patients undergoing radical cystectomy for aggressive bladder cancer in the national health service (NHS) and compared this with those with private health insurance (PHI).

Methods: This is retrospective study of 225 NHS and 32 PHI patients operated over 14 years by one surgeon. ASA scores were compared to approximate comorbidity status.

Results: There were significantly worse outcomes for all cause and disease specific mortality in the NHS group. There was no difference in ASA scores.

Conclusions: NHS patients fare worse than those with PHI. The reasons for this difference remain unclear. PHI patients have a more extensive lymphadenectomy and have fewer overall complications.

Keywords: NHS private insurance outcomes radical cystectomy bladder cancer

Introduction

We studied the outcome of patients with private health insurance (PHI) compared to national health service (NHS) patients. These patients had the same operation for muscle invasive cancer of the bladder, and we looked to see whether there were significant differences in both all-cause mortality and disease specific mortality. We have attempted to see whether we can propose any explanatory cause for such differences.

There are barriers in society that delay detection and the treatment of cancers for those people with various social and economic disadvantages [1]. This imbalance creates health disparities such as differences in survival [2]. There are racial, gender, ethnic and socioeconomic components which are reflected in their insurance status [3].

Previous studies, mostly in the United States, although with some English and Danish studies, have looked at the effect of

insurance status on outcomes of treatment for various cancers, including urological cancers of the prostate, kidney and bladder [4]. In the USA, the uninsured and those on Medicaid, have a worse stage at presentation [3-7] and a lower all cause survival than those with private insurance [2,1,8]. Worse disease specific mortality has also been shown to be an independent risk factor with worse outcomes for lower socio-economic status groups [9]. This finding, however, was not confirmed by a British study specifically looking at social deprivation [10]. Definitions of what constitutes adverse social groups are complex and may affect robustness of data [11]. A shift in classification of groups, pathological definitions and coding of cause of death may explain the lack of improvement in survival but not socioeconomic inequalities [11].

Many factors, including socio-demographic and clinical entities, form a complex network influencing survival. In the USA Insurance and socioeconomic status are predictors of mortality [1,8].

Uninsured patients are often treated at low volume centres[3]. Race and gender can also affect survival [12,13] as can communal poverty [14,15].

It has been suggested that poorer health generally, as demonstrated by more comorbidities and unhealthy lifestyle behaviours account for this phenomenon [2,15]. The affordable care act in the USA is an attempt to address the problem of the uninsured presenting with more advanced disease and their subsequent lower survival rates [16]. In addition, a negative association between income and prevalence of physicians for bladder and renal cancer mortality was found [17]. Those on Medicaid were less likely to receive definitive treatment [18]. Similar effects of socioeconomic status have been found in England [19]and Denmark [20].

These studies have all looked at the effect of all the range of treatments, both curative and palliative, for all types of cancer, including low- and high-grade disease which is of particular importance when discussing bladder cancer.

For differences in outcomes after specific surgical treatments, research has focused on radical prostatectomy. Insurance status is associated with stage and cancer control [21]. Higher surgery rates and lower radiotherapy rates were observed in private patients who were more affluent [22]. We have shown previously that insurance status is a protective risk factor with radical prostatectomy, but wealth itself is not protective which may be due to different adverse lifestyle behaviour[23].

There are few published studies pertaining to insurance status and survival after a specific treatment for bladder cancer, particularly radical cystectomy [24].

Effect of comorbidity and ASA score

Greater comorbidities as assessed by the ASA score, increase the likelihood of complications after radical cystectomy [25]. Patients with an ASA score of three or more have a greater complication rate [25]. Severe comorbidity is also associated with increased mortality [26,27]. The ASA score has been used to predict outcomes [28] and should be used for counselling and risk stratification [29,30]. In younger patients the ASA score can be useful despite EAU guidelines

suggesting otherwise [31,32]. Neglect of comorbidities can cause misleading impressions of results [33] but the ASA score can be useful at to ascertain complication risk particularly with the type of urinary diversion and the time taken to do more complex surgery [34]. Comorbidity assessed using the Charlson score is useful for younger and older patients to predict toxicity of treatment and outcomes [35,36]. All-cause mortality has a stronger relation to comorbidity than disease specific mortality does [37].

Urinarydiversion

In general, the neobladder has a higher complication rate and longer stay in hospital [38] although it may be offered to most patients [39-41]. However, in practice, it still tends to be reserved for younger and fitter patients and thus may serve as a proxy for generally better health [42].

Methods

This is a retrospective study. The 257 radical cystectomy with bilateral lymphadenectomy (with ileal conduit or neobladder) operations were performed by one surgeon in both NHS and the private sector in the same county of the UK, operated from 1999 to 2013 and followed up to the present day. There were 225 patients from the NHS and 32 patients with private health insurance. These were sequential patients who underwent potentially curative operative and neoadjuvant/adjuvant treatments.

We compared various patient characteristics from health records using Iqutopia for operation details, ICE, Bostwick laboratories and Masterlab for pathology and PACS for imaging.

Medical statistics were done using Medcalc software for logistic regression analysis to determine significant predictors of mortality and progression. Fishers, Mann-Whitney, t tests and log rank tests to compare Kaplan Meier. P values less than 0.05 were considered significant.

The ASA score was documented by the attending anaesthetist and retrieved from Iqutopia where available.

Results

(Table 1-4) (Graph1-6)

Table 1: Patient characteristics.

Characteristic	NHS n = 225	PHI n = 32	P
Mean age years	68 (IQR 62-76)	64 (IQR 59-71)	0.026
Mean tumour volume cc	23 (IQR 3-17)	11 (IQR 4-8.75)	0.18
Positive surgical margins	32 (14%)	2 (6%)	0.27
Stage localised	146 (56%)	20 (62%)	0.84
Locally advanced	99 (44%)	12 (38%)	
Carcinoma in situ	92 (41%)	19 (59%)	0.057
Mean number of nodes dissected	9 (IQR 12.2-14.2)	13 (IQR 6.0-18.0)	0.006
Patients with nodal involvement	43 (19%)	7 (22%)	0.811
Nodal extracapsular extension	24/43	7-Apr	1

Complications	89 (40%)	4 (13%)	0.002
Additional treatment given	38 (17%)	4 (13%)	0.79
Neobladder	40 (18%)	12 (38%)	0.16

Table 2: Significant predictors of mortality on logistic regression.


All-cause Mortality	Coefficient	P	Odds ratio
Neobladder	-0.8	0.0154	0.42
Private patient	-1.4	0.0013	0.24
Stage T3	1.007	0.0008	2.7
Stage T4	1.97	<0.0001	7.18
Disease Specific Mortality			
Additional treatment	0.77	0.021	2.17
Complication	0.73	0.013	2.08
Neobladder	-0.95	0.015	0.38
Node density	2.26	0.014	9.67
Private patient	-1.08	0.035	0.33
Progression			
Additional treatment	0.74	0.029	2.11
Private patient	-1.92	0.002	0.14
Stage T2	1.18	0.01	3.28
Stage T3	1.99	<0.0001	7.3
Stage T4	1.89	0.0001	6.6

Table 3: 1 year, 5 year and 10-year comparisons of mortality rates.


	NHS	PP
ACM 1 year	0.75	0.84
DSM 1 year	0.82	0.87
ACM 5 years	0.45	0.74
DSM 5 years	0.81	0.84
ACM 10 years	0.34	0.53
DSM 10 years	0.58	0.75

Table 4: ASA score and log rank tests.


	NHS	NHS	PP	PP	P value
	n = 151	%	18		
ASA 1	22	14	1	6	
ASA 2	95	63	12	67	
ASA 3	34	23	5	27	
ASA 1+2	117	77	13	73	0.56 Fishers 2 tailed
Mean ASA	2.1		2.2		0.52 Mann-Whitney
Log rank test	Chi squared	Degrees freedom	P value		
All-cause mortality NHS v PP	5.018	1	0.025		
Disease specific mortality NHS v PP	4.467	1	0.034		




Graph 1:




Graph 2:




Graph 3:



Graph 4:



Graph 5: All-cause mortality comparison



Graph 6: Disease specific mortality comparisons

There is a significant difference between the ages of the two populations with NHS patients being older by four years. Private patients had a greater degree of nodal dissection and fewer complications. However, there was no difference in tumour volume, positive surgical margin status, advanced stage, carcinoma in situ, nodal involvement, nodal extracapsular extension. Further, there was no difference in rates of neo adjuvant or adjuvant treatment or the use of a neobladder. Logistic regression showed private health status to have highly significant negative coefficients which are protective in all cause and disease specific mortality as well as lower progression rates.

All-cause mortality was associated with negative coefficients for neobladder and PHI. Locally advanced stage was predictive of a deleterious outcome.

Disease specific mortality also showed adverse coefficients for additional treatments (adjuvant and neoadjuvant), complications and nodal density involvement.

Progression was adversely affected by additional treatment and increasing stage.

The ASA score was available in 169 of the 257 patients (66%). There was no difference in the distributions of healthy and

unhealthy patients as measured by this score. (ASA1 and 2 were summated).

Log rank tests showed significant differences between the two groups for mortality.

Discussion

This study has supported generally what is known about bladder cancer and the negative impact of lower socio-economic status. However former studies have looked at bladder cancer as a whole, which includes low grade and non-muscle invasive disease as well as advanced muscle invasive disease, inoperable cases, cases treated with radiotherapy and palliative cases all grouped together. This study looks at a subsection of these, those deemed appropriate for radical cystectomy. We have seen significant differences in both all-cause mortality and disease specific mortality. These differences have not been previously demonstrated. There are many gross and subtle differences that are difficult to detect with such a heterogenous disease in a heterogenous population. We propose that looking at a subgroup like this will decrease some of the obfuscating factors, by limiting geographical variation and the operation being performed by one surgeon only.

All-cause mortality

We have detected a significant difference in all-cause mortality (Graph 5) with NHS patients (Graph 2) faring far worse (log rank $P = 0.025$) and with an odds ratio 0.24 in favour of PP (Graph 1, Table 2). Overall survival at one year is 0.75 for NHS and 0.84 for PHI, and at five years it is 0.45 for NHS and 0.74 for PHI. Similarly, at ten years for NHS patients it is 0.34 compared to 0.53 for PP (Table 3). Numerous explanations have been proposed for the worse prognosis overall for bladder cancer in the uninsured [1-12]. Overall worse health is the proposed principal reason for a worse all cause survival. We have tried to reflect potentially worse comorbidities with the ASA score as an approximation to overall health. There was, surprisingly, no significant difference detected between the two groups ($P = 0.56$) (Table 4). Private patients were four years younger on average and any chronic disease may not have reached a critical point to effect outcome ($P = 0.026$) (Table 1). This was not a significant factor on regression analysis.

We do see that having a neobladder is a significant ($P = 0.0154$) protective (negative coefficient -0.8) factor, perhaps standing as a proxy for overall health [38,39], but there was no significant difference in the use of this between the two groups ($P = 0.16$) (Table 1). Locally advanced stage 3 and 4 were both significant as expected [43, 44] with increasingly worse odds ratios (Table 2). The five-year gold standard for radical cystectomy is about 50% [43-49], and our population in the NHS reflects this. The patients with PHI fare much better with a 5-year survival of 0.74.

Disease specific mortality

We found a significant protective coefficient and odds ratio of 0.33 in favour of private patients (Table 2). There was a DSM of 0.87, 0.84 and 0.75 for PHI at one, five and ten years (Graph 3). This was superior to NHS patients with DSM of 0.82, 0.81 and 0.58 (Table 3, Graph 4). This compares favourably with other series (49). The log rank test comparing the two was significant (Graph 6, Table 5). Interestingly, the NHS mass of tumour was twice that of PHI, yet there was no significant difference in tumour volume which one might expect if there were a longer time to diagnosis in NHS patients (11cc v 23cc $P = 0.18$). Further, we did not detect a worse stage (NHS 56% localised, PHI 62% localised $P = 0.84$) or a significantly greater positive surgical margin rate in NHS patients (14% NHS v 6% PHI $P = 0.27$). However, the trend was worse with all of these parameters for NHS patients. There was no difference in rates of CIS between the two.

The use of neobladder was, once again, a protective risk factor (Table 2) which was used in 38% of PHI patients compared to 18% in NHS patients, however this did not reach significance ($P = 0.16$) (Table 1).

Additional treatments, neo-adjuvant and adjuvant therapies, were a significant detrimental risk factor (Table 2) but with no significant difference between the two groups (NHS 17% and PHI 13% $P = 0.79$).

The role of lymphadenectomy revealed no difference in severity of extracapsular invasion (Table 1) ($P = 1.0$) or in nodal involvement with 19% of NHS patients having involved metastatic deposits within at least one node, compared to 22% of PHI patients ($P = 0.811$). However, there was a significant difference in the number of nodes retrieved at lymphadenectomy, with an average of 9 for NHS and 13 for PHI patients ($P = 0.006$). Node density is a significant predictor of DSM with an odds ratio of 9.67 (Table 2).

Complications, of all types, are a significant adverse predictor for DSM. With a significantly worse rate in NHS, 40% compared to PHI patients, 13% ($P = 0.002$).

Progression

This also revealed PHI status to be protective and worse outcomes were associated with need for additional treatment and advancing stage.

Time to diagnosis and time to treatment from diagnosis may be a factor, although studies are conflicting [50,51]. There is no lead time bias (perceived increased survival with no effect on course of cancer) as there is no screening protocol.

Conclusions

American studies have shown that insurance is important and that even Medicaid patients do worse than private patients. We concur that in the UK the NHS does not provide equal outcomes compared with PHI when comparing radical cystectomy. There may be social infrastructure issues involving hospital building capacity, education, prevention, detection and treatment. Assistance to patients to access and navigate healthcare systems and workplace policies to encourage overall better health is needed. National policies have been implemented to tackle this complex problem. The NHS cancer plan 2000 in the United Kingdom (<http://www.doh.gov.uk/cancer/pdfs/cancerplan.pdf>) aims to reduce inequalities in survival between rich and poor with a 62-day target to treatment. The affordable care act in the USA highlights benefits of extending health coverage to the uninsured [1].

We have found significant differences in outcomes for all cause and disease specific mortality, yet the cause is elusive. We have shown a direct comparison of NHS and PHI to reveal these outcomes which can remain hidden in American studies as the SEER database does not include uninsured patients by definition. There is no difference in tumour volume, carcinoma in situ, advanced stage or not, nodal involvement, extracapsular involvement of nodes, rates of adjuvant and neo-adjuvant treatment. There are, however, differences in complication rates and extent of lymphadenectomy with favourable features pertaining to PHI.

We need to look for other ways to measure overall health that include medical and lifestyle with socioeconomic components.

Limitations

The PHI cohort is small. We have not measured the time to diagnosis and to treatment. We have not been able to document the Charlson comorbidity score. We have not documented types of complications.

References

1. Ward E, Halpern M, Schrag N, Cokkinides V, DeSantis C, et al. (2008) Association of insurance with cancer care utilization and outcomes. *CA Cancer J Clin* 58(1): 9-31.
2. Niu X, Roche LM, Pawlish KS, Henry KA (2013) Cancer survival disparities by health insurance status. *Cancer Med* 2(3): 403-411.
3. Jacobs BL, Montgomery JS, Zhang Y, Skolarus TA, Weizer AZ, et al. (2012) Disparities in bladder cancer. *Urol Oncol* 30(1): 81-88.
4. Ward EM, Fedewa SA, Cokkinides V, Virgo K (2010) The association of insurance and stage at diagnosis among patients aged 55 to 74 years in the national cancer database. *Cancer J* 16(6): 614-621.
5. Halpern MT, Ward EM, Pavluck AL, Schrag NM, Bian J, et al. (2008) Association of insurance status and ethnicity with cancer stage at diagnosis for 12 cancer sites: a retrospective analysis. *Lancet Oncol* 9(3): 222-231.
6. Urinary Bladder Cancer in Florida - Bureau of Epidemiology.
7. Shackley DC, Clarke NW (2005) Impact of socioeconomic status on bladder cancer outcome. *Curr Opin Urol* 15(5): 328-331.
8. Du XL, Lin CC, Johnson NJ, Altekruse S (2011) Effects of individual-level socioeconomic factors on racial disparities in cancer treatment and survival: Findings from the National Longitudinal Mortality Study, 1979-2003. *Cancer* 117(14): 3242-3251.
9. Siddiqui MM, Heney NM, McDougal WS, Feldman AS (2015) Disparities in overall and urothelial carcinoma specific mortality associated with healthcare insurance status. *Bladder* 2(1): e10.
10. Begum G, Dunn JA, Bryan RT, Bathers S, Wallace DM (2004) Socio-economic deprivation and survival in bladder cancer. *BJU Int* 94(4):539-543.
11. A Shah, B Rachet, E Mitry, N Cooper, CM Brown, et al. (2008) Survival from bladder cancer in England and Wales up to 2001. *Br J Cancer* 99(Suppl 1): S86-S89.
12. Mallin K, David KA, Carroll PR, Milowsky MI, Nanus DM (2011) Transitional cell carcinoma of the bladder: racial and gender disparities in survival (1993 to 2002), stage and grade (1993 to 2007). *J Urol* 185(5):1631-1636.
13. A Moran, AM Sowerbutts, S Collins, N Clarke, R Cowan (2004) Bladder cancer: Worse survival in women from deprived areas. *British Journal of Cancer* 90: 2142-2144.
14. Brookfield KF, Cheung MC, Gomez C, Yang R, Nieder AM (2009) Survival disparities among African American women with invasive bladder cancer in Florida. *Cancer* 115(18): 4196-4209.
15. Kaye DR, Canner JK, Kates M, Schoenberg MP, Bivalacqua TJ (2016) Do African American Patients Treated with Radical Cystectomy for Bladder Cancer have Worse Overall Survival? Accounting for Pathologic Staging and Patient Demographics Beyond Race Makes a Difference. *Bladder Cancer* 2(2): 225-234.
16. Mahal BA, Aizer AA, Ziehr DR, Hyatt AS, Lago-Hernandez C (2014) The association between insurance status and prostate cancer outcomes: implications for the Affordable Care Act. *Prostate Cancer Prostatic Dis* 17(3): 273-279.
17. Janet Colli, Leah Grossman, Oliver Sartor, Benjamin R Lee (2011) 50 Urologic cancer mortality rates stratified by geographic region and physician prevalence in the united states. *Journal of Urology General & Epidemiological Trends & Socioeconomics: Evidence-based Medicine & Outcomes* 185(4s): e21-e22.
18. Mahal AR, Mahal BA, Nguyen PL, Yu JB (2018) Prostate cancer outcomes for men aged younger than 65 years with Medicaid versus private insurance. *Cancer* 124(4): 752-759.
19. MP Coleman, B Rachet, LM Woods, E Mitry, M Riga, et al (2004) Trends and socioeconomic inequalities in cancer survival in England and Wales up to 2001. *British Journal of Cancer* 90(7):1367-1373.
20. Kirsten Thorup Eriksen, Anne Petersen, Aslak Harbo Poulsen, Isabelle Deltour, Ole Raaschou-Nielsen (2008) Social inequality and incidence of and survival from cancers of the kidney and urinary bladder in a population-based study in Denmark, 1994-2003. *European Journal of Cancer* 44(14):2030-2042.
21. Andrea Gallina, Pierre I Karakiewicz, Felix K-H Chun, Alberto Briganti, Hartwig Huland, et al. (2007) Health-insurance status is a determinant of the stage at presentation and of cancer control in European men treated with radical prostatectomy for clinically localized prostate cancer. *BJU Int* 99(6): 1404-1408.
22. Barbiere JM, Greenberg DC, Wright KA, Brown CH, Palmer C, et al. (2012) The association of diagnosis in the private or NHS sector on prostate cancer stage and treatment. *J Public Health* 34(1): 108-114.
23. Simon Robinson, Marc Laniado, Assad Farooq, Hanif Motiwala, Mohamed Omar, et al. A comparison of clinical parameters at presentation, pathological outcomes and biochemical relapse between NHS and private patients undergoing radical prostatectomy at a single centre in the United Kingdom. *Journal of clinical urology* 9(5): 326-334.
24. Robinson S, Rao A (2015) Differences in mortality and patient characteristics between national health service patients and those with

- private health insurance post radical cystectomy. Resource Centre - EMUC15.
25. Novotny V, Zastrow S, Koch R, Wirth MP (2012) Radical cystectomy in patients over 70 years of age: impact of comorbidity on perioperative morbidity and mortality. *World J Urol* 30(6): 769-776.
 26. Fairey A, Chetner M, Metcalfe J, Moore R, Todd G, et al. (2008) Associations among age, comorbidity and clinical outcomes after radical cystectomy: results from the Alberta Urology Institute radical cystectomy database. *J Urol* 180(1): 128-134.
 27. Bartosz Dybowski, Krzysztof Ossoliński, Anna Ossolińska, Michal Peller, EwaBres-Niewada, et al. (2015) Impact of stage and comorbidities on five-year survival after radical cystectomy in Poland: single centre experience. *Cent European J Urol* 68(3): 278-283.
 28. Aziz A, May M, Burger M, Palisaar RJ, Trinh QD, et al. (2014) Prediction of 90-day mortality after radical cystectomy for bladder cancer in a prospective European multicenter cohort. *Eur Urol* 66(1): 156-163.
 29. Boorjian SA, Kim SP, Tollefson MK, Carrasco A, Chevillie JC, et al. (2013) Comparative performance of comorbidity indices for estimating perioperative and 5-year all-cause mortality following radical cystectomy for bladder cancer. *J Urol* 190(1): 55-60.
 30. Williams SB, Kamat AM, Chamie K, Froehner M, Wirth MP, et al. (2018) Systematic Review of Comorbidity and Competing-risks Assessments for Bladder Cancer Patients. *Eur Urol Oncol* 1(2): 91-100.
 31. Froehner M, Koch R, Hübler M, Heberling U, Novotny V, et al. (2018) Predicting 90-day and long-term mortality in octogenarians undergoing radical cystectomy. *BMC Urol* 18(1): 91.
 32. Bokarica P, Hrkac A, Gilja I (2017) Re: J. Alfred Witjes, Thierry Lebret, Eva M. Compérat, et al. Updated 2016 EAU Guidelines on Muscle-invasive and Metastatic Bladder Cancer. *Eur Urol* 2017; 71: 462-75. *Eur Urol* 72(2): e45.
 33. Feinstein AR (1970) The pre-therapeutic classification of co-morbidity in chronic disease. *J Chronic Dis* 23(7): 455-468.
 34. Malavaud B, Vaessen C, Mouzin M, Rischmann P, Sarramon J, et al. (2001) Complications for radical cystectomy. Impact of the American Society of Anesthesiologists score. *Eur Urol* 39(1): 79-84.
 35. Extermann M, Overcash J, Lyman GH, Parr J, Balducci L (1998) Comorbidity and functional status are independent in older cancer patients. *J Clin Oncol* 16(4):1582-1587.
 36. Figueroa AJ, Stein JP, Dickinson M, Skinner EC, Thangathurai D, et al. (1998) Radical cystectomy for elderly patients with bladder carcinoma: an updated experience with 404 patients. *Cancer* 83(1):141-147.
 37. Miller DC, Taub DA, Dunn RL, Montie JE, Wei JT (2003) The impact of co-morbid disease on cancer control and survival following radical cystectomy. *J Urol* 169(1): 105-109.
 38. Evans B, Montie JE, Gilbert SM (2010) Incontinent or continent urinary diversion: how to make the right choice. *Curr OpinUrol* 20(5): 421-425.
 39. Autorino R, Quarto G, Di Lorenzo G, De Sio M, Perdonà S, et al. (2009) Health related quality of life after radical cystectomy: comparison of ileal conduit to continent orthotopic neobladder. *Eur J Surg Oncol* 35(8): 858-864.
 40. Lawrentschuk N, Colombo R, Hakenberg OW, Lerner SP, Månsson W, et al. (2010) Prevention and management of complications following radical cystectomy for bladder cancer. *Eur Urol* 57(6): 983-1001.
 41. Lee KS, Montie JE, Dunn RL, Lee CT (2003) Hautmann and Studer orthotopic neobladders: a contemporary experience. *J Urol* 169(6): 2188-2191.
 42. Saika T, Suyama B, Murata T, Manabe D, Kurashige T, et al. (2001) Orthotopic neobladder reconstruction in elderly bladder cancer patients. *Int J Urol* 8(10): 533-538.
 43. Bassi P, Ferrante GD, Piazza N, Spinadin R, Carando R, et al. (1999) Prognostic factors of outcome after radical cystectomy for bladder cancer: a retrospective study of a homogeneous patient cohort. *J Urol* 161(5): 1494-1497.
 44. Dalbagni G, Genega E, Hashibe M, Zhang ZF, Russo P, et al. (2001) Cystectomy for bladder cancer: a contemporary series. *J Urol* 165(4): 1111-1116.
 45. Shariat SF, Karakiewicz PI, Palapattu GS, Lotan Y, Rogers CG, et al. (2006) Outcomes of radical cystectomy for transitional cell carcinoma of the bladder: a contemporary series from the Bladder Cancer Research Consortium. *J Urol* 176(6 Pt 1): 2414-2422.
 46. Ghoneim MA, el-Mekresh MM, el-Baz MA, el-Attar IA, Ashamallah A (1997) Radical cystectomy for carcinoma of the bladder: critical evaluation of the results in 1,026 cases. *J Urol* 158(2): 393-399.
 47. Stein JP, Skinner DG (2006) Radical cystectomy for invasive bladder cancer: long-term results of a standard procedure. *World J Urol* 24(3): 296-304.
 48. Williams SB, Huo J, Chu Y, Baillargeon JG, Daskivich T, et al. (2017) Cancer and All-cause Mortality in Bladder Cancer Patients Undergoing Radical Cystectomy: Development and Validation of a Nomogram for Treatment Decision-making. *Urology* 110: 76-83.
 49. Taekmin Kwon, In Gab Jeong, Dalsan You, Bumsik Hong, Jun Hyuk Hong, et al. (2014) Long-Term Oncologic Outcomes after Radical Cystectomy for Bladder Cancer at a Single Institution. *J Korean Med Sci* 29(5): 669-675.
 50. Nielsen ME1, Palapattu GS, Karakiewicz PI, Lotan Y, Bastian PJ, et al. (2007) A delay in radical cystectomy of >3 months is not associated with a worse clinical outcome. *BJU Int* 100(5): 1015-1020.
 51. Gore JL, Lai J, Setodji CM, Litwin MS, Saigal CS (2009) Mortality increases when radical cystectomy is delayed more than 12 weeks: Results from a Surveillance, Epidemiology, and End Results-Medicare analysis. *Cancer* 115(5): 988-996.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)

DOI: 10.32474/JUNS.2020.03.000151



Journal of Urology & Nephrology Studies

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles